P19771.P01,

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No.

P19771

Total Pages

Inventor(s) or Application Identifier Susumu SEINO, Tadao SHIBASAKI, and Nobuaki OZAKI

Title: PROTEIN RIM2

ADDRESS TO:

Assistant Commissioner for Patents Box Patent Application Washington, DC 20231

	S	APPLICATION ELEMENTS	ACCOMPANYING APPLICATION PARTS
	- Piro	Fee Transmittal Form Specification [Total Pages 41]	8. Assignment Papers (cover sheet & document(s))
		- Descriptive title of the Invention - Cross References to Related Applications - Statement Regarding Fed sponsored R & D - Reference to Microfiche Appendix - Background of the Invention - Brief Summary of the Invention - Brief Description of the Drawings (if filed) - Detailed Description - Claim(s) - Abstract of the Disclosure	9. □ 37 CFR 3.73(b) Statement (when there is an assignee) 10. □ English Translation Document (if applicable) 11. □ Information Disclosure Statement (IDS)/PTO-1449 12. ☒ Preliminary Amendment 13. ☒ Return Receipt Postcard (MPEP 503) (Should be specifically itemized)
3.		Drawing(s) (35 USC 113) [Total Sheets 8]	14. ⊠ Small Entity ☐ Statement filed in prior application.
4.5.6.7.	a. b.		Statements(two) Status still proper and desired 15. The prior application is assigned of record to 16. Foreign priority claimed
18.		Ta CONTINUING APPLICATION, check appropriate box and sup Continuation	ply the requisite information: of prior Application No/, filed
19.		Amend the specification by inserting before the first line the sentence. This application is a continuation-in-part, continuation, di	ce: vision, of Application No/, filed

Address all future correspondence to Customer No. 7055 at the present address of:

GREENBLUM & BERNSTEIN, P.L.C.

1941 Roland Clarke Place
Reston, VA 20191
(703) 716-1191

Bruce H. Bernstein, Reg No. 29,027 Typed or Printed Name

[Page 1 of 1]

GREENBLUM & BERNSTEIN, P.L.C.

PATENT AND TRADEMARK CAUSES 1941 ROLAND CLARKE PLACE RESTON, VIRGINIA 20191

Applicant or Patentee: Susumu SEINO et al.	Serial or Patent No:
Attorney's Docket No.:	
Filed or Issued: For: Protein Rim2	_
101, A 2000M Mille	
VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY	
STATUS (37 CFR 1.9 (f) and 1.27(c)) SMALL BUSINESS CONCERN	
I hereby declare that I am	
the owner of the small business concern identified below:	•
[V] an official of the small business concern empowered to act on behalf of the concern identified below:	
NAME OF SMALL BUSINESS CONCERN JCR Pharmaceuticals Co., Ltd.	
ADDRESS OF SMALL BUSINESS CONCERN 3-19 Kasuga-cho, Ashiya-shi, Hyogo 659-0021 Japan	
I hereby declare that the above identified small business concern qualifies as a small business concern as defined in 13 CFR 121.12 and repro of paying reduced fees to the United States Patent and Trademark Office, in that the number of employees of the concern, including thos persons. for purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other vaconcern controls or has the power to control the other, or a third party or parties controls or has the power to control both.	e of its affiliates, does not exceed 500 f the concern of the persons employed
I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with contract Protein Rim2 by inventor(s) Susumu SEINO et al. described in	n regard to the invention,
[] the specification filed herewith	
[] the specification filed herewith [] application serial no	
[] patent no, issued	
Pube rights held by the above identified small business concern are not exclusive, each individual, concern or organization having rights in the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9 (a), or a nonprofit organization under 37 CFR 1.9 (b), any concern which would not qualify as a small business concern under 37 CFR 1.9 (d), or a nonprofit organization under 37 CFR 1.9 (e), or a nonprofit organization under 37 CFR 1.9 (e) from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 (e) [1.5] [(c) if that person made the invention, *NOTE: Separate verified statements
ADDRESS 22-1-4 Aobanomorinomachi, 638-1 Chibateramachi, Chuo-ku, Chiba-shi, Chiba 260-0844 Japan	-
[/] INDIVIDUAL [] SMALL BUSINESS CONCERN [] NONPROFIT ORGANIZATION	•
FULL NAME	
4DDRESS	-
[] INDIVIDUAL [] SMALL BUSINESS CONCERN [] NONPROFIT ORGANIZATION	
acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate.	y status prior to paying, or at the time (37 CFR 1.28 (b))
hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are beliet attements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any p s directed.	under section 1001 of Title 18 of the
NAME OF PERSON SIGNING Shin Ashida	
TITLE OF PERSON IF OTHER THAN OWNER President	
ADDRESS OF PERSON SIGNING <u>JCR Pharmaceuticals Co., Ltd., 3-19 Kasuga-cho, Ashiya-shi, Hyogo 659-0021</u>	Japan
SIGNATURE June 27, 2000	

NF301.WPF - SES FORM SMALL BUSINESS CONCERN

GREENBLUM & BERNSTEIN, P.L.C. PATENT AND TRADEMARK CAUSES

PATENT AND TRADEMARK CAUSES 1941 ROLAND CLARKE PLACE RESTON, VIRGINIA 20191

Applicant or Patentee:	Susumu SEINO et al.	
Serial or Patent No:	Attorney's Docket No.:	
Filed or Issued: For: Protein]	Rim2	
101		
	VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS (37 CFR 1.9 (f) and 1.27(b)) — INDEPENDENT INVENTOR	
	entor, I hereby declare that I qualify as an independent inventor as defined in 37 CFR 1.9 (c) for purposes of paying reduced fees alted States Code, to the Patent and Trademark Office with regard to the invention entitled	
described in		
[[] the specification filed herewith [] application serial no, filed	
who could not be classi	anted, conveyed or licensed and am under no obligation under contract or law to assign, grant, convey or license, any rights in the invisified as an independent inventor under 37 CFR 1.9 (c) if that person had made the invention, or to any concern which would not 37 CFR 1.9 (d) or a nonprofit organization under 37 CFR 1.9 (e).	ention to any person of qualify as a small
	or organization to which I have assigned, granted, conveyed, or licensed or am under an obligation under contract or law to assigne invention is listed below:	ın, grant, convey, or
fi.	[] no such person, concern, or organization [✓] persons, concerns or organizations listed below*	
	*NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)	
FULL NAMEJ	JCR Pharmaceuticals Co., Ltd.	
ADDRESS3-19	JCR Pharmaceuticals Co., Ltd. 19 Kasuga-cho, Ashiya-shi, Hyogo 659-0021 Japan 10 ASMALL BUSINESS CONCERN L. INONPROFIT ORGANIZATION	
[] INDIVIDUA	AL [1/] SMALL BUSINESS CONCERN [] NONPROFIT ORGANIZATION	
FULL NAME		
ADDRESS		
[] INDIVIDUA	AL [] SMALL BUSINESS CONCERN [] NONPROFIT ORGANIZATION	
ADDRESS	AL [] SMALL BUSINESS CONCERN [] NONPROFIT ORGANIZATION	
= [] INDIVIDOA	AL [] SWALL BUSINESS CONCERN [] MONTROTTI ON GAMEATION	
I acknowledge the duty to time of paying, the earlie	to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior liest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFF	to paying, or at the R 1.28 (b))
these statements were ma	statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be trade with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under so Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to	ection 1001 of Title
Susumu SEIN		
NAME OF INVEN	NTOR NAME OF INVENTOR NAME OF INVENTOR	
Musus A	flow Tadao Shihasahi Nobraki Ozaki	
Signature of Inventor	Signature of Inventor Signature of Inventor	
Jun 29,	2000 June 29, 2000 June 28, 2000	
Date	Date Date	

(NF300.WPF - SES FORM INDEPENDENT INVENTOR)

P19771.A01

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant : Susumu SEINO et al.

Serial No : Not Yet Assigned

Filed : Concurrently Herewith

For : PROTEIN RIM2

PRELIMINARY AMENDMENT

Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

Prior to calculation of the filing fees and the examination of the above-identified patent application on the merits, the Examiner is respectfully requested to amend the claims as follows:

IN THE CLAIMS

Please amend the claims as follows:

Claim 3, line 1, delete "or 2"

Claim 5, lines 3 and 4, delete "or 2"

Claim 10, line 2, change "one of claims 4 to 7" to ---claim 4---.

Claim 13, line 2, delete "or 2"

REMARKS

By the above amendment, the claims have been amended to delete multiple dependency.

If there should be any questions, the Examiner is invited to contact the undersigned at the telephone number listed below.

Respectfully submitted, Susumu SEINO et al.

Bruce H. Bernstein

Reg. No. 29,027

July 14, 2000 GREENBLUM & BERNSTEIN, P.L.C. 1941 Roland Clarke Place Reston, VA 20191 (703) 716-1191

10

15

20

25

30

PROTEIN RIM2

FIELD OF THE INVENTION

The present invention relates to protein Rim2, which is a novel isoform of Rim, i.e., a protein that interacts with a low molecular G protein Rab3 and is proposed to serve as a regulator of Rab3-dependent synaptic vesicle fusion, and which specifically interacts with the GDP/GTP exchange factor (GEFII; a cAMP sensor). More specifically, the present invention relates to elucidation of the mechanisms of intracellular vesicle transport and secretion, and to the novel protein Rim2 which is useful in diagnosis of endocrine-related diseases or neuropathy and in development of agents for prevention and treatment thereof, the gene encoding Rim2 and an antibody addressed to Rim2 protein.

Rim2 is considered to be a regulatory factor of vesicle fusion. It was found in the course of the present invention that the protein is expressed predominantly in endocrine tissues and endocrine- and neuroendocrine-derived cell lines. GTP-Rab3/GEFII/Rim complex is thought to participate in the regulation of exocytosis of neurons and endocrine cells, in a cAMP-dependent and protein kinase A (PKA) independent manner.

BACKGROUND OF THE INVENTION

Transport of substances between cell organelles, which are unit membrane-enclosed structures such as endoplasmic reticulum, is conducted by intracellular vesicle transport. In endocrine cells including pancreatic β -cells and pituitary cells, peptides/proteins synthesized at ribosomes are received by the endoplasmic reticulum, from which they are transported in vesicles, which are transformed into secretory vesicles through the Golgi body and transported to the cell membrane, where they are released out of the cell via a step which includes fusion of the membranes. In neurons, neurotransmitter-containing precursors of synaptic vesicles are formed in Golgi bodies and transported by microtubules along the axon and stored at the synapse. Depolarization of the pre-synaptic membrane causes the vesicles to fuse with the pre-synaptic membrane and thus the neurotransmitters are released. This type of secretion based on the fusion of the vesicles and the cell membrane is called exocytosis.

In contrast, when extracellular substances such as hormones including cell

in it is a second of the secon

5

10

15

20

25

30

growth factors are bound to the cell membrane, the complexes thus formed are invaginated into the cell to form endosomes. This type of uptake of environmental substances is called endocytosis.

Formation of vesicles, such as by budding, commonly observed both in exocytosis and endocytosis, and docking and fusion, the phenomena observed in process of their transportation and binding to other membrane systems, are regulated by a GTP-binding, low-molecular protein, called G protein. More than 30 types of this protein are known. The group of the proteins, which are also classified in Rab family, regulate the intracellular vesicle transport system.

With regard to the intracellular vesicle transport system, it is understood today that a cell is in a resting state when Rab protein occurs in a bound form to guanine nucleotide diphosphate (GDP), and that budding, docking and fusion are triggered as a result of a process in which a protein having GEF activity act on Rab protein and converts it to GTP-binding Rab protein, to which GTP binds to form a GTP-Rab complex, which in turn binds to a corresponding target protein on the membrane.

Stimulus-secretion coupling plays an important role in exocytosis observed in many cell types including neurons and endocrine cells [J.E. Rothman, Nature 372:55(1994); T.C. Sudhof, Nature 375:645 (1995)]. While a rise in intracellular Ca²⁺ concentration is important in the regulation of exocytosis, other signals are play important roles. cAMP (cyclic adenosine-3',5'also known to monophosphate)/PKA (cAMP-dependent protein kinase A) signaling pathway is known to regulate exocytosis in many of neurons, neuroendocrine cells and endocrine cells. In particular, cAMP has been thought to mediate long-term potentiation by increasing neurotransmitter release in the brain [R.D. Hawkins et al. Ann. Rev. Neurosci. 16:625(1993); G. Lonart et al., Neuron 21:1141(1998)]. cAMP also regulates exocytosis responsible for insulin release from pancreatic β cells and amylase release from parotid acinar cells [P.M. Jones and S.J., Persaud, Endocrine. Rev. 19:429(1998); E. Renstrom, et al., J. Physiol. 502:105(1997); K. Yoshimura, Biochim. Biophys. Acta 1402:171(1998)].

In addition to its role in PKA-dependent phosphorylation of regulatory proteins associated with the process of exocytosis, it is known that cAMP also acts directly on the exocytotic machinery in neurons and non-neuronal cells [G. Lonart et al., Neuron 21:1141 (1998); E. Renstrom et al., J. Physiol. 502:105 (1997); K.

Yoshimura, Biochim. Biophys. Acta, 1402:171(1998)].

During the search by the yeast two-hybrid screen (i.e., a method for detection of the interaction between two proteins in yeast cells) for an intracellular signaling molecule directly coupling to a sulphonylurea receptor, a component of pancreatic β -cell ATP-sensitive K⁺ (K_{ATP}) channels [N. Inagaki et al. Proc. Natl. Acad. Sci. U.S.A. 91,2679 (1994)], a cAMP sensor protein (called "CAMPS") was identified and it was found that the protein has two putative cAMP binding domains, a Pleckstrin homology domain (PH domain), and a guanine nucleotide exchange factor (GEF) homology domain.

In the course of this study, two study groups independently reported cAMP binding proteins that activate Rap1, a member of the small G binding proteins [J. de Rooiji et al. Nature 396:474 (1998); H. Kawasaki et al. Science 282:2275 (1998)], and CAMPS was incidentally revealed to be a mouse homologue of cAMP-GEFII [H. Kawasaki et al. Science 282:2275 (1998)].

Though the mechanisms of intracellular vesicle transport system have thus gradually been clarified, substantial part of them remains still unknown. Further progress is needed for the understanding of the mechanisms so as to provide diagnostic agents or therapeutics for a variety of diseases which involve neurons or endocrine cells.

Unlike the former suggestion that only a single cAMP binding domain was present in cAMP-GEFII, the study by the present inventors suggested the presence of two putative cAMP binding domains (cAMP-A and cAMP-B), based on a sequence alignment of cAMP-GEFII sequence and regulatory subunits of PKA. Figure 1 shows the sequence alignment of the cAMP binding domains. The cAMP binding domains A and B (cAMP-A and cAMP-B, respectively) of cAMP-GEFII and the cAMP binding domains A and B of the PKA regulatory subunit I α (RI α -A and RI α -B, respectively) are shown. The invariant residues in the different cAMP-binding domains are indicated by black boxes.

As shown in Figure 2, a glutathione-S-transferase (GST)-cAMP-A fusion protein bound to [3 H]cAMP with a dissociation constant (Kd) of $^{-10}\mu$ M, while the binding of [3 H]cAMP to a GST-cAMP-B fusion protein was not evident under the same conditions.

Figure 2 shows the binding of cAMP to cAMP-A. GST-cAMP-A (filled circles) or GST-PKA RI α (open circles) was incubated with different concentrations

20

25

5

10

15

of [3 H]cAMP (0-50 μ M). The data for cAMP-A or PKA RI α are normalized relative to maximal cAMP binding activities. Kd values are $10.0\pm2.3~\mu$ M and 23.7 ± 0.6 nM for cAMP and PKA RI α , respectively.

In the cAMP-B domain, the amino acid residue 423, which originally is glutamic acid (Glu), is substituted with lysine (Lys). This glutamic acid residue is important for cAMP binding. Considering that a more rapid dissociation than the wild-type was observed with a PKA regulatory subunit having an equivalent mutation (E-200-K), cAMP-B may also dissociate cAMP rapidly. Thus, a possibility remains that cAMP binds to the cAMP-B domain.

10

15

20

25

SUMMARY OF THE INVENTION

As identification of a target molecule of CAMPUS, cAMP-GEFII, would serve to show its physiological role, the present inventors attempted to find a molecule that interacts with cAMP-GEFII by means of a yeast two-hybrid screen (YTH) method on the MIN6 cDNA library (See "Identification of Interacting molecules by YTH Method").

Surprisingly, the present inventors found that cAMP-GEFII interacts with a novel isoform (named "Rim2" by the present inventors) of Rim (a molecule which specifically interacts with Rab3: Rab3-interacting molecule: Hereinafter referred to as "Rim1"). Rim1 protein is a putative effector of the small G protein Rab3 and is proposed to serve as a Rab3-dependent regulator of synaptic vesicle fusion [Y. Wang et al. Nature 388:593(1997)].

The full-length novel protein Rim2 sequenced by the present inventors, which consists of 1590 amino acid residues, was found to have 61.6 % identity with rat Rim1. As Figure 3 shows, a zinc finger, PDZ and two C2 domains were found highly conserved between Rim1 and Rim2.

Based on the above findings, the present invention provides a protein having the amino acid sequence set forth under SEQ ID NO:1 in the Sequence Listing.

30

The present invention further provides a protein having an amino acid sequence with one or more amino acids deleted, substituted, inserted or added relative to the amino acid sequence set forth under SEQ ID NO:1 in the Sequence Listing and which has a property to interact with GDP/GTP exchange factor II.

The present invention further provides a mouse gene which encodes the

15

20

25

30

following proteins (1) or (2):

- (1) a protein having the amino acid sequence set forth under SEQ ID NO:1 in the Sequence Listing,
- (2) a protein having an amino acid sequence with one or more amino acids deleted, substituted, inserted or added relative to the above-identified amino acid sequence and which has a property to interact with GDP/GTP exchange factor II.

In the present specification, "one or more" amino acid residues are generally several (e.g., 3 or 4) to 10 residues.

The present invention further provides a DNA having a nucleotide sequence set forth under SEQ ID NO:2 in the Sequence Listing, the DNA being a cDNA corresponding to the above protein having the amino acid sequence set forth under SEQ ID NO:1 in the Sequence Listing.

The present invention further provides a DNA having a nucleotide sequence with one or more nucleotides deleted, substituted, inserted or added relative to the nucleotide sequence set forth under SEQ ID NO:2 in the Sequence Listing and encoding any one of the above proteins. Herein, "one or more" nucleotides are generally several (e.g., 3 or 4) to 10 nucleotides. A variety of such nucleotide sequences with one or more nucleotides deleted, substituted, inserted or added can be readily prepared by those skilled in the art by making use of the familiar knowledge on degeneracy of the genetic code.

The present invention further provides a DNA having the nucleotide sequence of the coding region of the any one of the above DNA's or of a DNA having the nucleotide sequence set forth under SEQ ID NO:2 in the Sequence Listing.

The present invention further provides a DNA fragment consisting of a part of any one of the above DNA's.

The present invention further provides a probe comprising a DNA which hybridizes with the DNA consisting of any one of the above nucleotide sequences.

The present invention further provides a primer DNA fragment consisting of a partial sequence of any one of the above nucleotide sequences.

The present invention further provides a recombinant vector having any one of the above DNA's.

The present invention further provides a monoclonal or polyclonal antibody directed to any one of the above proteins.

The present invention further provides a diagnostic agent for human use

10

15

20

25

30

comprising any one of the above probes or antibodies. The diagnostic agent is useful in the test for such diseases as secretion disorders in secretory systems including pituitary, hypothalamus, pancreatic β -cells and parotid gland, or the test for brain-nervous system diseases.

The present invention further provides a therapeutic agent for any one of the above diseases.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 illustrates a sequence alignment of the cAMP binding domains.

Figure 2 is a graph showing the binding of cAMP to cAMP-A.

Figure 3 illustrates a comparison of amino acid identity between Rim1 and Rim2, in zinc finger, PDZ and C2 domains.

Figure 4 shows the results of immunoblotting showing the interaction between cAMP-GEFII and Rim1 or Rim2.

Figure 5 shows the results of Northern blot analysis of cAMP-GEFII, Rim1 and Rim2 in various rat tissues and endocrine- and neuroendocrine-derived cell lines.

Figure 6 is the result of *In situ* hybridization showing the localization of Rim1 and Rim2 in mouse brain and pituitary.

Figure 7 is a graph showing the result of yeast two-hybrid assays.

Figure 8 illustrates the result of immunoblotting showing the interaction between Rab3A and Rim1 or Rim2 in vitro.

Figure 9 is a graph showing the time course for high K*-induced GH secretion from PC12 cells cotransfected with GH and cAMP-GEFII.

Figure 10 is a graph showing the effect of forskolin on GH secretion from transfected PC cells.

Figure 11 is a graph showing forskolin-induced GH secretion from PC12 cells transfected with various mutant cAMP-GEFII.

Figure 12 is a graph showing the effect of H-89 on forskolin-induced GH secretion from PC12 cells transfected with cAMP-GEFII.

Figure 13 is a schematic illustration showing a model for cAMP-dependent exocytosis.

DETAILED DESCRIPTION OF THE INVENTION

10

15

20

25

30

A variety of mutants can be provided by means of recombinant DNA technology. First, mutations can be introduced into a DNA clone fragment through different chemical and/or enzymatic processes, and the mutant DNA's thus obtained are then sequenced to select particular mutants with intended merits. This method allows a systematic preparation of different mutants regardless of their phenotypes. General methods of preparing a mutant clone DNA are as follows.

- 1. With the help of an oligonucleotide, substitution, deletion, insertion or addition can be directly effected in a given DNA sequence. This method enables to introduce a number of mutations in a small region of a given DNA.
- 2. By using longer oligonucleotides, it is possible to synthesize a desired gene.
- 3. By means of region-specific mutagenesis, a desired mutation can be introduced into a large (1-3 kb) DNA region.
- 4. Linker-scanning mutagenesis of DNA is a method suited for introducing a cluster point mutation into a relatively small (4-10 bp) DNA region.
- 5. PCR is also utilized as a method for direct introduction of a mutation. [References: Current Protocols in Molecular Biology., 3 Vols., Edited by Ausubel F.M. et al., John Wiley & Sons, Inc., Current Protocols., Vol. 1, Chapter 8: Mutagenesis of Cloned DNA, pages 8.0.1-8.5.10]

Also well known to those skilled in the art are methods of preparing plasmids or vectors which can express a desired gene including different mutations obtained by the above methods. That is, by inserting a DNA carrying a desired gene into a expression vector DNA using a combination of restriction enzymes and a ligase, a recombinant plasmid is readily constructed which carries the desired gene. The recombinant plasmid thus obtained is then introduced into different cells to transfect them, thereby producing transformed cells. Cells which may be utilized range from prokaryotes, e.g. *E. coli*, to yeast, insect, plant and animal cells. [References: Vectors Essential Data. Gacesa P. and Ramji D.P., 166 pages. BIOS Scientific Publishers Limited 1994., John Wiley & Sons in association with BIOS Scientific Publishers Ltd. Expression vectors, pages 9-12.]

Introduction of a recombinant plasmid into host cells is effected by calcium chloride method or electroporation. Calcium chloride method provides efficient transformation without requiring any special apparatus. For higher efficiency,

10

15

20

25

30

electroporation is recommended.

[References: Current Protocols in Molecular Biology, 3 Vols. Edited by Ausbel F.M. et al., John Wiley & Sons, Inc., Current Protocols, Vol. 1, unit 1.8: Introduction of Plasmid DNA into Cells, pages 1.8.1-1.8.10]

Two types are known of transfection generally carried out on animal cell lines, i.e., transient and permanent types. In transient transfection, transformed cells are cultured for 1 - 4 days to effect transcription and replication of the transfected gene, and then the cells are harvested and their DNA analyzed. Alternatively, in many studies, a stable transformant cell line is produced, in which the transfected gene is incorporated into the chromosomes. Examples of the method for transfection include calcium phosphate method, electroporation, and liposome fusion method.

[Reference: Current protocols in molecular biology. 3 vols. Edited by Ausubel F.M. et al., John Wiley & Son, Inc., Current Protocols. Vol. 1, chapter 9: Introduction of DNA into mammalian cells, pages 9.0.1-9.17.3.]

Polyclonal and monoclonal antibodies directed to the proteins (polypeptides) coded by Rim2 gene of the present invention or their fragments and analogues as well, are readily prepared using techniques well known in the art. Antibodies obtained may be used as laboratory reagents and diagnostic agents for diseases associated with Rim2 gene. The antibodies obtained are also used for preparation of antibody columns, for immunoprecipitation as well as for identification of the antigen by Western blotting.

A general method for preparing a monoclonal antibody in mg-scale directed to the proteins coded for by Rim2 gene of the present invention is as follows: Mice are inoculated with the antigen protein to immunize. The spleen is removed from the mice exhibiting a sufficient antibody titer. The spleen cells are dissociated, and selected B cells are fused with myeloma cells of B cell origin to form hybridoma cells which secrete the antibody. The monoclonal antibody secreted from the hybridoma cells is purified from the culture medium using an affinity column, ion-exchange, or gel filtration, etc. The polyclonal antibody of the present invention may be prepared by a conventional method: Using rabbits, horses, mice or guinea pigs as immunized animals, the antigen protein is inoculated along one of the schedules known in the art to immunize the animals, and then IgG, etc. are isolated from the collected serum.

15

20

25

30

[Reference: Current protocols in molecular biology, 3 vols. Edited by Ausubel F.M. et al., John Wiley & Sons, Inc., Current Protocols, Vol. 2, chapter 11: Immunology, pages 11.0.1-11.16.13.]

In order to assess the specificity of interaction between cAMP-GEFII and Rim2, the present inventors evaluated the binding of a FLAG-tagged cAMP-GEFII protein to a GST-Rim2 fusion protein immobilized on glutathione beads (See "Study on Interaction between Rim2 and cAMP-GEFII: I").

Briefly, lysates from COS-1 cells transfected with FLAG-tagged cAMP-GEFII, from MIN6 cells or from mouse brain homogenate were evaluated for binding to GST-Rim1, GST-Rim2 or GST alone. cAMP was detected by immunoblotting with an anti-FLAG antibody (Figure 4, left) or an anti-cAMP-GEFII antibody (Figure 4, center and right), respectively. These results demonstrates that cAMP-GEFII protein interacts with GST-Rim2 protein. Likewise, GST-Rim1 protein also bound to cAMP-GEFII in the mouse brain homogenate (See "Study on Interaction between Rim1 and cAMP-GEFII") (Figure 4, right). These results confirms that cAMP-GEFII interacts with Rim1 and Rim2.

Figure 5 shows the results of the northern blot analyses of cAMP-GEFII, Rim1 and Rim2 in various rat tissues and endocrine- and neuroendocrine-derived cell lines (See "Northern Blotting in Rat Tissues"). Ten μ g samples of total RNA from various tissues and cell lines (except 5 μ g for pancreatic islets) were used. Hybridization and washing were performed under standard conditions. The faint signals seen in Rim2 mRNA blot analysis of cerebrum and cerebellum are due to cross-hybridization with the Rim1 cDNA probe used. Figure 5 shows that Rim2 mRNA is expressed predominantly in endocrine tissues and endocrine- and neuroendocrine-derived cell lines, including pituitary, pancreatic Langerhans' islet cells, MIN6 cells, and PC12 cells. Rim2 mRNA was detected in the brain by reverse transcriptase-PCR (data not shown). Rim1 mRNA, in contrast, was found to be expressed in cerebrum, cerebellum, and pituitary by a similar analysis.

The major transcripts for Rim1 and Rim2 have 6.4 kb for Rim1, and 7.2 kb and 5.4 kb for Rim2. There are also found several minor transcripts, which occur due probably to alternative splicing.

cAMP-GEFII mRNA is generally coexpressed with Rim1 or Rim2 mRNA in tissues and cell lines in which regulated exocytosis is known to occur. Figure 6 illustrates the results of *in situ* hybridization showing the localization of Rim1 and

10

15

20

25

30

Rim2 in mouse brain and pituitary. In the figure: (a) cAMP-GEFII; (b) Rim1; (c) Rim2; (d) pituitary. The scale bar corresponds to 1 mm. Abbreviations: Cb = cerebellum, Cp = caudoputamen, Cx = cortex, Hi = hippocampus, Ob = olfactory bulb, Po = pons, Th = thalamus

Rim2 mRNA is found expressed only in the cerebellar cortex, while Rim1 mRNA is expressed in cerebral cortex, hippocampus (especially CA3 and dentate gyrus), olfactory bulb, and cerebellar cortex (See "In situ Hybridization in Mouse Brain"). The distribution of cAMP-GEFII mRNA overlaps largely with that of Rim1 mRNA in the brain. It is confirmed that Rim2 mRNA and cAMP-GEFII mRNA are coexpressed in anterior pituitary.

Rim1 is proposed to be a Rab3 effector, a low molecular weight G protein [Y. Wang, et al., Nature 388,593 (1997)]. Using yeast two-hybrid assays (See "Study on Interaction between Rim2 and Rab3A".), the present inventors found that Rim2, like Rim1, interacts with active Rab3A (Q81L) (Figure 7). Figure 7 shows the results of the yeast two-hybrid assays. Rim1, Rim2 or rabphilin3 and wild-type Rab3A or constitutively active Rab3A (Q81L) in various combinations were determined by transactivation of liquid β -galactosidase activity.

In addition, the immobilized GST-Rim2 bound only to the GTP γ S-bound form of Rab3A (Figure 8). Figure 8 shows the interaction between Rab3A and Rim1 or Rim2 in vitro, which is the result obtained by incubating GTP γ S- or GDP γ S-bound form of Rab3A with GST-Rim1 (residues 1-201) and GST-Rim2 (residues 1-345) immobilized on glutathione beads, respectively. Rab3A was detected by immunoblotting with anti-Rab3A antibody. These results indicate that Rim2, like Rim1, binds to the GTP-activated form of Rab3A.

The interaction of cAMP-GEFII and Rim2 protein strongly suggests that cAMP-GEFII is involved in regulated exocytosis. To determine its functional role, the present inventors examined the effect of cAMP on Ca2⁺-dependent secretion in PC12 cells cotransfected with growth hormone (GH) and cAMP-GEFII (See "Study on GH secretion from Transfected PC12 Cells").

Since PC12 cells endogenously express Rim2 but not cAMP-GEFII, the exogenously introduced cAMP-GEFII may form a complex with endogenous Rim2.

Figure 9 is a graph showing the time course of high K⁺-induced GH secretion from PC12 cells cotransfected with GH and cAMP-GEFII. Figure 10 is a graph showing the effect of forskolin on GH secretion from the transfected PC12

10

15

20

25

30

cells. Forskolin (50 μ M) was added 10 min before the incubation with a low K⁺ (4.7 mM) or high K⁺ (60 mM) solution. The meaning of the symbols are as follows: For basal (low K⁺-induced) secretion: cAMP-GEFII-transfectant (filled triangles); β -galactosidase-transfectant (control)(open circles); high K⁺-induced secretion: cAMP-GEFII-transfectant (filled circles); β -galactosidase-transfectant (control)(open circles). The values represent the percent GH amounts released into the medium relative to the total cellular GH amounts.

In the cotransfected PC12 cells, as shown in Figure 9, cAMP-GEFII did not alter Ca²⁺-dependent (60 mM K⁺) secretion of cotransfected GH, compared to the control, but significantly enhanced forskolin (50 μ M)-induced, Ca²⁺-dependent GH secretion (Figure 10). Forskolin acts mainly on adenylate cyclase, serving to increase cAMP levels in the cells. cAMP-GEFII also enhanced 8-Br-cAMP (1 mM)-induced, Ca²⁺-dependent GH secretion (cAMP-GEFII-transfectant, 34.9 \pm 1.3 %; control, 25.1 \pm 1.8 %, n=9, P<0.001).

Figure 11 is a graph showing forskolin-induced GH secretion from PC12 cells transfected with various mutant cAMP-GEFII, in which the increment of forskolin (50 μ M)-induced GH secretion (in the presence of high K⁺) above the basal level during a 15-min incubation for each mutant cAMP-GEFII is expressed as percentage relative to the wild-type cAMP-GEFII (100%). In the figure: WT = wild-type cAMP-GEFII, T810A = mutant cAMP-GEFII (T810A); G114E, G422D = double mutant cAMP-GEFII (G114, G422D).

The forskolin-induced GH secretion was not affected in the mutant cAMP-GEFII (T810A) in which a potential PKA phosphorylation site is disrupted by substitution of one of its amino acids (Figure 11). In addition, the forskolin-induced GH secretion in the mutant cAMP-GEFII (G114E, G422D) in which both of the cAMP binding sites are disputed was reduced to ~40 % of that in the wild-type.

These results indicate that cAMP promotes Ca²⁺-dependent GH secretion by binding to cAMP-GEFII, without involving its phosphorylation by PKA.

Figure 12 is a graph showing the effect of H-89 on forskolin-induced GH secretion from cAMP-GEFII-transfected PC12 cells. H-89 (10 μ M) was added to the incubation buffer 10 min before forskolin (50 μ M) treatment. The treatment with H-89 (10 μ M) reduced high K⁺-induced GH secretion in both of the cAMP-GEFII-transfected and β -galactosidase-transfected PC12 cells. The data were obtained from 3-5 independent experiments (A-D). The values are means \pm SEM

(P<0.01).

5

10

15

20

25

30

Importantly, the forskolin-induced, Ca2*-dependent GH secretion from the cAMP-GEFII-transfected PC12 cells treated with the PKA inhibitor H-89 was significantly higher than that from the control cells. This indicates that cAMP-GEFII mediates cAMP-dependent and PKA-independent exocytosis.

To ascertain the physiological relevance of cAMP-GEFII, the present inventors investigated the role of endogenous cAMP-GEFII in secretion. In insulin secretion from pancreatic β -cells, cAMP is proposed to stimulate exocytosis by PKA-dependent as well as PKA-independent mechanisms [M. Prentki, F.M. Matschinsky, Physiol. Rev. 67:1185 (1987)/ P.M.Jones, S.J. Persaud, Endocrine. Rev. 19:429 (1998)].

In the high glucose condition of 16.7 mM, 8-Br-cAMP-induced insulin secretion from MIN6 cells treated with antisense oligonucleotides against cAMP-GEFII was significantly reduced (87.5 ± 2.3 % of the secretion from MIN6 cells treated with a control oligonucleotide, n=27, P<0.005) (See "Study of the Role of cAMP-GEFII in cAMP-dependent Exocytosis"), suggesting that cAMP-GEFII participates in cAMP-dependent exocytosis in native cells.

Rab3 is associated with the final step of exocytosis. The structurally-related proteins rabphilin3 [H. Shirataki et al., Mol. Cell. Biol. 13,2061 (1993)] and Rim1 both bind to Rab3A, suggesting that multiple Rab3A effectors could operate in triggering docking and fusion of the vesicles to the plasma membrane.

In the process toward the present invention, it was found that the cAMP sensor, cAMP-GEFII, mediates cAMP-induced, Ca²⁺-dependent exocytosis by interacting with a Rab3 effector Rim2.

In addition to its role in PKA phosphorylation of proteins associated with secretory processes, previous studies have suggested that cAMP may act directly on the exocytosis [G. Lonart, et al., Neuron 21:1141 (1998); E. Renstrom, et al., J. Physiol. 502:105(1997); K. Yoshimura et al., Biochim. Biophys. Acta 1402:171(1998)]. In pancreatic β -cells, too, PKA-dependent as well as PKA-independent stimulation of insulin release by cAMP has been proposed [E. Renstrom, et al., J. Physiol. 502:105 (1997)]. It is thought that cAMP probably directly stimulates amylase release in parotid acinar cells [G. Lonart, et al., Neuron 21:1141 (1998)]. In addition, a recent study suggests that cAMP enhances glutamate release in the brain partly by a direct action on the exocytotic machinery

[G. Lonart, et al., Neuron 21,1141 (1998)].

However, while both rabphilin3 and Rim1 are ubiquitously expressed in most of the synapses in the brain[C. Li et al., Neuron 13:885 (1994)], cAMP-enhanced glutamate release occurs in synaptosomes from the CA3 region in the hippocampus, not from the CA1 region, a finding consistent with cAMP-GEFII and Rim1 being coexpressed predominantly in CA3.

Accordingly, it is considered that, in addition to PKA-dependent phosphorylation in the secretory processes, cAMP promotes regulated exocytosis in a PKA-independent manner by acting directly on a complex of cAMP-GEFII (a cAMP sensor) and Rim (a Rab3 effector) in some neurons and neuroendocrine and endocrine cells, as schematically illustrated in Figure 13.

These findings indicates that Rim2 of the present invention also plays an important role in the regulation of exocytosis in neurons and endocrine cells.

15 EXAMPLES

5

10

20

25

30

The present invention will be described below in further detail by presenting specific procedures in the present invention with reference to an example.

<Sequencing of CAMPS (cAMP-GEFII) cDNA>

A plasmid cDNA library has been made from a mouse insulin-secreting cell line, MIN6, in the vector pVP16. A yeast two-hybrid bait vector was constructed in plasmid pBTM116 using a DNA fragment encoding partial rat SUR1 (amino acid residues 598-1003)(GenBank accession number L40624), a subunit of the pancreatic β -cell K_{ATP} channel.

Yeast two-hybrid screen of the plasmid MIN6 cDNA library was performed as described in K. Kotake et al., J. Biol. Chem. 272:29407 (1997). A prey clone encoding a partial CAMPS, a cAMP sensor, (residues 187-730) was isolated. A full-length mouse CAMPS cDNA was obtained from the λ MIN6 cDNA library [N. Inagaki et al., Proc. Natl. Acad. Sci. U.S.A. 91:2679(1994)]. The nucleotide sequence of mouse CAMPS (cAMP-GEFII) has been deposited in Genbank with the accession number of AB021132.

<Preparation and Test of GST fusion Protein>

cAMP-A (amino acid residues 43-153), cAMP-B (amino acid residues 357-469), and rat PKA regulatory subunit (RI α)(full-length) were expressed as GST-

10

15

20

25

30

fusion proteins using pGEX-4T-1 (Amersham-Pharmacia) and purified according to the manufacturer's instructions. cAMP binding assay was performed as described in R.A. Steiberg, et al., J. Biol. Chem. 262:2664(1987) with slight modifications.

Briefly, GST-fusion protein (1 μ g) was incubated in binding buffer (200 μ l) containing various concentrations of [³H]cAMP, 50 mM potassium phosphate buffer (pH 6.8), 150 mM NaCl, 1 mM EDTA, 5 mM 2-mercaptoethanol, and 0.5 mg/ml bovine serum albumin with or without 40 mM unlabeled cAMP for 2 hrs on ice.

<Identification of Interacting molecules by YTH Method>

Yeast two-hybrid bait vector was constructed in plasmid pBTM116 using a full-length mouse cAMP-GEFII cDNA. A prey clone encoding a partial sequence of Rim2 (amino acid residues 53-863) was isolated from the plasmid MIN6 cDNA library. A full-length cDNA for Rim2 was obtained from the λ MIN6 cDNA library. <Study on Interaction between Rim2 and cAMP-GEFII: I>

Rim2 (amino acid residues 538-863) was expressed as a GST fusion protein and purified according to the method described in "Preparation and Test of GST fusion Protein". A full-length cAMP-GEFII cDNA was subcloned into plasmid pFLAG-CMV-2 (Sigma). The resultant construct was transfected into COS-1 cells, using LipofectAMINE (Life Technologies). The lysate of the COS-1 cells was incubated with GST-Rim2 immobilized on glutathione beads for 2 hrs at 4°C. The complex thus obtained was washed with distilled water, separated by SDS-PAGE, and immunoblotted with an anti-FLAG M2 antibody (Sigma).

<Study on Interaction between Rim2 and cAMP-GEFII: II>

The lysate of MIN6 cells was incubated with GST-Rim2 and interaction between cAMP-GEFII and Rim2 was evaluated according to the method described in "Study on Interaction between Rim2 and cAMP-GEFII: I", using a IgG antibody raised against the C-terminus (amino acid residues 1001-1011, Gln-Met-Ser-His-Arg-Leu-Glu-Pro-Arg-Arg-Pro) (SEQ ID NO:5) of mouse cAMP-GEFII.

<Study on Interaction between Rim1 and cAMP-GEFII>

According to the method described in "Preparation and Test of GST fusion Protein", Rim1 partial sequence (530-806) was expressed as a GST fusion protein and then purified. The brain homogenate from three mice was incubated with GST-Rim1 immobilized on glutathione beads overnight at 4°C. cAMP-GEFII was detected as described in "Study on Interaction between Rim2 and cAMP-GEFII: II".

<Northern Blotting in Rat Tissues>

Northern Blotting was performed for various tissues of rat using, as probes, mouse cAMP-GEFII (nucleic acids 606-2237), rat Rim1 (1035-1491), and mouse Rim2 (586-1490) cDNA.

5 < In situ Hybridization in Mouse Brain>

In situ hybridization in mouse brain was performed as described in J. Tanaka, M. Murate, C.Z. Wang, S. Seino, T. Iwanaga, Arch. Histol. Cytol. 59:485 (1996).

Antisense oligonucleotide probes (45 mer) used for mouse cAMP-GEFII and Rim2 correspond to the regions of the nucleic acids 2746-2790 and 1376-1420, respectively.

For the antisense oligonucleotide for Rim1, Rim1 cDNA was partially cloned from mouse brain: the probe used in this was 5'-ttgcgctcactcttctggcctcccttgccattctgctctgaaagc-3' (SEQ ID NO:3).

<Study on Interaction between Rim2 and Rab3A>

According to the method described in "Identification of Interacting molecules by YTH Method", the full-length cDNA's for wild type mouse Rab3A and constitutively active bovine Rab3A (Q81L) were cloned into the yeast bait vector pBTM116.

The nucleotide sequence of zinc finger domains of bovine rabphilin3 (amino acid residues 1-283), rat Rim1 (amino acid residues 1-204) and mouse Rim2 (amino acid residues 1-345 were cloned into the prey vector pVP16. Liquid culture assay of β -galactosidase activities was performed according to the manufacturer's instructions (Clontech). The activity values were obtained from 3 independent clones for each transformant and normalized by cell numbers determined as OD_{600} .

Lipid-modified Rab3A was purified from the membrane fraction of Sf9 cells expressing Rab3A. Rat Rim1 (amino acid residues 1-204) and mouse Rim2 (amino acid residues 1-345) were expressed as GST fusion proteins and purified. The GTP γ S- or GDP β S-bound form of Rab3A was incubated for 90 min at 4°C with GST-Rim1, or GST-Rim2 (30 pmol for each) immobilized on glutathione beads in reaction buffer. Rab3A was detected by immunoblotting with anti-Rab3A antibody. <Study on GH secretion from Transfected PC12 Cells>

GH secretion from transfected PC12 cells was performed as described in K. Korake et al., J. Biol. Chem., 272:29407(1997). Expression plasmid vectors (pSR

20

25

30

10

 α) for wild-type cAMP-GEFII, mutant cAMP-GEFII (T810A), and the double mutant cAMP-GEFII (G114E, G422D) were prepared. As a control, β -galactosidase (β -gal) was used. PC cells were transfected with GH expression vector (pXGH5: Nichols Institute) plus each vector described above, using LipofectAMINE.

PC12 cell were incubated with a low K⁺ (4.7 mM) or high K⁺ (60 mM) solution, in the presence or absence of forskolin (50 μ M) or 8-bromoadenosine 3',5' cyclic monophosphate (8-Br-cAMP)(1 mM). Forskolin or 8-Br-cAMP was added 10 min before the incubation with a low or high K⁺ solution. In some experiments, the PKA inhibitor H-89 (10 μ M) was added 10 min before forskolin stimulation.

<Study of the Role of cAMP-GEFII in cAMP-dependent Exocytosis>

To interfere with the synthesis of cAMP-GEFII in MIN6 cells, antisense phosphorothicate-substituted oligoDNA (16 mer) against mouse cAMP-GEFII (the region corresponding to nucleic acids 104-119) and control oligoDNA (5'-acctacgtgactacgt-3') (SEQ ID NO:4) were synthesized (BIOGNOSTIK).

MIN6 cells were treated with 4 μ M of the antisense oligoDNA or control oligoDNA 24 hours before insulin secretion experiments. The efficacy of antisense oligoDNA was evaluated by immunoblot analysis of the antisense oligoDNA-treated MIN6 cells over-expressing cAMP-GEFII by transient transfection, using anti-cAMP-GEFII antibody. The level of cAMP-GEFII was markedly lowered in the antisense oligoDNA-treated MIN6 cells. Insulin secretory response to 8-Br-cAMP (1 mM) of these MIN6 cells was assessed in the presence of high glucose (16.7 mM). Five separate experiments were performed, in which insulin was measured as described in T. Gonoi et al., J. Biol. Chem. 269:16989 (1994).

5

10

15

SEQUENCE LISTING

<110> Seino, Susumu; JCR Pharmaceuticals Co., Ltd.

5 <120> Protein Rim2

<130> GP35

<160>4

10 <210> 1

<211> 1590

<212> PRT

<213> Mus musculus

15 <400> 1

20

30

Met Ser Ala Pro Leu Gly Pro Arg Gly Arg Pro Ala Pro Thr Pro Ala 1 5 10 15

Ala Ser Gln Pro Pro Pro Gln Pro Glu Met Pro Asp Leu Ser His Leu 20 25 30

Thr Glu Glu Glu Arg Lys Ile Ile Leu Ala Val Met Asp Arg Gln Lys 35 40 45

25 Lys Glu Glu Glu Lys Glu Gln Ser Val Leu Lys Ile Lys Glu Glu His
50 55 60

Lys Ala Gln Pro Thr Gln Trp Phe Pro Phe Ser Gly Ile Thr Glu Leu 65 70 75 80

Val Asn Asn Val Leu Gln Pro Gln Gln Lys Gln Pro Asn Glu Lys Glu 85 90 95

Pro Gln Thr Lys Leu His Gln Gln Phe Glu Met Tyr Lys Glu Gln Val

Hard Could also the street of the street of

285

	Lys 1	Lys	Met 115	Gly	Glu	Glu	Ser	Gln 120	Gln	Gln	Gln	Glu	Gln 125	Lys	Gly	Asp
5	Ala	Pro 130	Thr	Cys	Gly	Ile	Cys 135	His	Lys	Thr	Lys	Phe 140	Ala	Asp	Gly	Cys
10	Gly 145	His	Asn	Cys	Ser	Tyr 150	Cys	Gln	Thr	Lys	Phe 155	Cys	Ala	Arg	Cys	Gly 160
	Gly	Arg	Val	Ser	Leu 165	Arg	Ser	Asn	Lys	Val 170	Met	Trp	Val	Cys	Asn 175	Leu
15	Cys	Arg	Lys	Gln 180	Gln	Glu	Ile	Leu	Thr 185	Lys	Ser	Gly	Ala	Trp 190	Phe	Tyr
	Asn	Ser	Gly 195	Ser	Asn	Thr	Leu	Gln 200	Gln	Pro	Asp	Gln	Lys 205	Val	Pro	Arg
20	Gly	Leu 210	Arg	Asn	Glu	Glu	Ala 215		Gln	Glu	Lys	Lys 220		Lys	Leu	His
25	G1u 225	Gln	Pro	Gln	Phe	G1n 230		Ala	Pro	Gly	Asp 235	Leu	Ser	Val	Pro	Ala 240
	Val	Glu	ı Lys	Gly	Arg 245		His	Gly	Leu	Thr 250		Gln	ı Asp	Thr	Ile 255	Lys
30	Asn	Gly	y Ser	Gly 260		Lys	His	s Gln	11e 265		ser	Asp	Met	Pro 270		Asp
	Arg	Ly	s Arg	g Sei	r Pro	Se 1	r Va	l Sei	r Arg	g Ası	Glr	ı Ası	n Arg	g Arg	g Tyr	Glu

280

	Gln	Ser 290	Glu	Glu	Arg	Glu	Asp 295	Tyr	Ser	Gln	Tyr	Val 300	Pro	Ser	Asp	Gly
5	Thr 305	Met	Pro	Arg	Ser	Pro 310	Ser	Asp	Tyr	Ala	Asp 315	Arg	Arg	Ser	Gln	Arg 320
	Glu	Pro	Gln	Phe	Tyr 325	Glu	Glu	Pro	Gly	His 330	Leu	Asn	Tyr	Arg	Asp 335	Ser
10	Asn	Arg	Arg	Gly 340	His	Arg	His	Ser	Lys 345	Glu	Tyr	Ile	Val	Asp 350	Asp	Glu
15	Asp	Val	Glu 355	Ser	Arg	Asp	Glu	Tyr 360	Glu	Arg	Gln	Arg	Arg 365	Glu	Glu	Glu
	Tyr	Gln 370	Ala	Arg	Tyr	Arg	Ser 375	Asp	Pro	Asn	Leu	Ala 380	Arg	Tyr	Pro	Val
20	Lys 385		Gln	Pro	Tyr	G1u 390	Glu	Gln	Met	Arg	Ile 395	His	Ala	Glu	Val	Ser 400
0.5	Arg	Ala	Arg	His	Glu 405		Arg	His	Ser	Asp 410	Val	Ser	Leu	Ala	Asn 415	Ala
25	Glu	Leu	Glu	Asp 420		Arg	Ile	Ser	Leu 425		Arg	Met	Asp	Arg 430	Pro	Ser
30	Arg	g Gln	1 Arg 435		· Val	Ser	Glu	Arg 440		: Ala	ı Ala	Met	Glu 445		Gln	Arg
	Sei	Tyı 45(. Met	Glu	ı Arg	Thr 455		: Glu	ı Ala	ı Glm	Gly 460		Ser	Ser	Tyr

	Pro 465	Gln	Arg	Thr	Ser	Asn 470	His	Ser	Pro	Pro	Thr 475	Pro	Arg	Arg	Ser	Pro 480
5	Ile	Pro	Leu		Arg 485	Pro	Asp	Met	Arg	Arg 490	Ala	Asp	Ser	Leu	Arg 495	Lys
	Gln	His	His	Leu 500	Asp	Pro	Ser	Ser	Ala 505	Val	Arg	Lys	Thr	Lys 510	Arg	Glu
10	Lys	Met	Glu 515	Thr	Met	Leu	Arg	Asn 520	Asp	Ser	Leu	Ser	Ser 525	Asp	Gln	Ser
	Glu	Ser 530	Val	Arg	Pro	Pro	Pro 535	Pro	Arg	Pro	His	Lys 540	Ser	Lys	Lys	Gly
15	Gly 545	Lys	Met	Arg	Gln	Val 550	Ser	Leu	Ser	Ser	Ser 555	Glu	Glu	Glu	Leu	Ala 560
20	Ser	Thr	Pro	Glu	Tyr 565	Thr	Ser	Cys	Asp	Asp 570		Glu	Leu	Glu	Ser 575	Glu
	Ser	Val	Ser	Glu 580		Gly	Asp	Ser	Gln 585		Gly	Lys	Arg	Lys 590		Ser
25	Glu	Glm	Gly 595		Leu	ı Ser	Asp	Ser 600		Thr	Arg	Ser	Glu 605		Gln	Lys
	Lys	610		. Tyr	· Tyr	: Gly	Gly 615		Ser	· Leu	ı Glu	Glu 620		Leu	Glu	Trp
30	Se 1		ı Pro	Gln	ı Ile	e Lys 630		Sei	Gly	v Val	l Asp 635		· Cys	s Ser	· Ser	Thr 640
	Th	r Le	u Ası	ı Glı	ı Glı	u His	s Ser	His	s Sei	r Ası	p Lys	His	s Pro	o Val	Thr	Trp

۳	Gln	Pro	Ser	Lys 660	Asp	Gly	Asp	Arg	Leu 665	He	Gly	Arg	Ile	Leu 670	Leu	Asn
5	Lys	Arg	Leu 675	Lys	Asp	Gly	Ser	Va1 680	Pro	Arg	Asp	Ser	Gly 685	Ala	Met	Leu
10	Gly	Leu 690	Lys	Val	Val	Gly	Gly 695	Lys	Met	Thr	Glu	Ser 700	Gly	Arg	Leu	Cys
	Ala 705	Phe	Ile	Thr	Lys	Val 710	Lys	Lys	Gly	Ser	Leu 715	Ala	Asp	Thr	Val	Gly 720
15	His	Leu	Arg	Pro	Gly 725	Asp	Glu	Val	Leu	G1u 730	Trp	Asn	Gly	Arg	Leu 735	Leu
20	Gln	Gly	Ala	Thr 740	Phe	Glu	Glu	Val	Tyr 745	Asn	Ile	Ile	Leu	GI u 750	Ser	Lys
20	Pro	Glu	Pro 755	Gln	Val	Glu	Leu	Val 760	Val	Ser	Arg	Pro	11e 765	Gly	Asp	Ile
25	Pro	Arg 770	He	Pro	Asp	Ser	Thr 775	His	Ala	Gln	Leu	Glu 780	Ser	Ser	Ser	Ser
	Ser 785	Phe	Glu	Ser	Gln	Lys 790	Met	Asp	Arg	Pro	Ser 795	Ile	Ser	Val	Thr	Ser 800
30	Pro	Met	Ser	Pro	Gly 805	Met	Leu	Arg	Asp	Val 810	Pro	Gln	Phe	Leu	Ser 815	Gly
	Gln	Leu	Ser	Ile	Lys	Leu	Trp	Phe	Asp	Lys	Val	Gly	His	Gln	Leu	Ile

	Val	Thr	11e 835	Leu	Gly	Ala	Lys	Asp 840	Leu	Pro	Ser	Arg	Glu 845	Asp	Gly	Arg
5	Pro	Arg 850	Asn	Pro	Tyr	Val	Lys 855	He	Tyr	Phe	Leu	Pro 860	Asp	Arg	Ser	Asp
10	Lys 865	Asn	Lys	Arg	Arg	Thr 870	Lys	Thr	Val	Lys	Lys 875	Thr	Leu	Glu	Pro	Lys 880
10	Trp	Asn	Gln	Thr	Phe 885	Ile	Tyr	Ser	Pro	Val 890	His	Arg	Arg	Glu	Phe 895	Arg
15	Glu	Arg	Met	Leu 900	Glu	Ile	Thr	Leu	Trp 905	Asp	Gln	Ala	Arg	Val 910	Arg	Glu
	Glu	Glu	Ser 915	Glu	Phe	Leu	Gly	Glu 920	He	Leu	Ile	Glu	Leu 925	Glu	Thr	Ala
20	Leu	Leu 930	Asp	Asp	Glu	Pro	His 935	Trp	Tyr	Lys	Leu	G1n 940	Thr	His	Asp	Val
O.E.	Ser 945	Ser	Leu	Pro	Leu	Pro 950	Arg	Pro	Ser	Pro	Tyr 955	Leu	Pro	Arg	Arg	Gln 960
25	Leu	His	Gly	Glu	Ser 965	Pro	Thr	Arg	Arg	Leu 970	Gln	Arg	Ser	Lys	Arg 975	He
30				980					985		Glu			990		
	Val	Ser	Asp 995	Tyr	Arg	His	Asn	Gly 1000		Asp	Leu	GIN	Ser 100		Inr	Leu

	Ser	Val 1010		Glu	Gln	Val	Met 1015		Ser	Asn	His	Cys 1020		Pro	Ser	Gly
5	Ser 1025		His	Arg	Val	Asp 103(Ile	Gly	Arg	Thr 1035		Ser	Trp	Ser	Pro 1040
	Ser	Ala	Pro	Pro	Pro 1045		Arg	Asn	Val	Glu 1050		Gly	His	Arg	Gly 1058	
10	Arg	Ala	Thr	Gly 1060		Tyr	Asn	Thr	11e 1065		Arg	Met	Asp	Arg 1070		Arg
15	Val	Met	Asp 1075		His	Tyr	Ser	Ser 1080		Arg	Asp	Arg	Asp 1085	Cys	Glu	Ala
10	Ala	Asp 1090		Gln	Pro	Tyr	His 1095		Ser	Arg	Ser	Thr 1100		Gln	Arg	Pro
20	Leu 110		Glu	Arg	Thr	Thr 1110		Arg	Ser	Arg	Ser 1119		Glu	Arg	Pro	Asp 1120
	Thr	Asn	Leu	Met	Arg 1125		Met	Pro	Ser	Leu 1130		Thr	Gly	Arg	Ser 1135	
25	Pro	Pro	Ser	Pro 1140		Leu	Ser	Arg	Ser 1145		Pro	Arg	Thr	Gly 1150		Val
20	Gln	Thr	Ser 115		Ser	Ser	Thr	Pro 1160		Thr	Gly	Arg	Arg	Gly 5	Arg	Gln
30	Leu	Pro		Leu	Pro	Pro	Lys		Thr	Leu	Glu	Arg		Ala	Met	Asp
	Ile	Glu	Glu	Arg	Asn	Arg	Gln	Met	Lys	Leu	Asn	Lys	Tyr	Lys	Gln	Val

E	Ala	Gly	Ser	Asp	Pro 1205		Leu	Glu	Gln	Asp 1210		His	Ser	Lys	Tyr 1215	
5	Ser	Gly	Trp	Asp 1220		His	Arg	Gly	Ala 1225		Thr	Val	Ser	Thr 1230		Ser
10	Ser	Asp	Ser 123	Asp	Val	Ser	Asp	Val 1240		Ala	Val	Ser	Arg 1245		Ser	Ser
	Ala	Ser 125		Phe	Ser	Ser	Thr 1259		Tyr	Met	Ser	Val 1260		Ser	Glu	Arg
15	Pro 126		Gly	Asn	Arg	Lys 1270		Ser	Val	Phe	Thr 1275		Lys	Met	Gln	Asn 1280
	Arg	Gln	Met	Gly	Val 128		Gly	Lys	Asn	Leu 1290		Lys	Ser	Thr	Ser 129	
20	Ser	Gly	Asp	Met		Ser	Leu	Glu	Lys 130		Asp	Gly	Ser	Gln 131		Asp
25	Thr	Ala	Va1 131	Gly 5	Ala	Leu	Gly	Thr 132		Gly	Lys	Lys	Arg		Ser	Ser
	Ile	Gly 133		. Lys	Met	Val	Ala 133		Val	Gly	Leu	Ser 134		Lys	Ser	Arg
30	Ser 134		Ser	· Gln	Leu	Ser 135		Thr	Glu	Gly	Gly 135		Lys	Lys	Leu	Arg 1360
	Ser	Th 1	· Val	Gln	Arg 136		Thr	Glu	Thr	Gly 137		Ala	Val	Glu	Met 137	

	Asn	Trp	Me t	Thr	Arg	Gln	Ala	Ser	Arg	Glu	Ser	Thr	Asp	Gly	Ser	Met
				1380	1				1385					1390		
5	Asn	Ser	Tyr 1395		Ser	Glu	Gly	Asn 1400	Leu)	Ile	Phe	Pro	Gly 1405		Arg	Leu
	Ala	Ser 1410		Ser	Gln	Phe	Ser 1415		Phe	Leu	Asp	Gly 1420		Gly	Pro	Ala
10	Gln 1425	Leu		Gly	Arg	Gln 1430	Thr		Ala	Thr	Pro 1435	Ala		Gly	Asp	Ile 1440
15			Gly	Met	Met 144	Asp		Lys	Gly	Gln 1450	Leu		Val	Glu	I le 1455	Ile
	Arg	Ala	Arg	Gly 1460	Leu		Val	Lys	Pro 146	Gly		Lys	Thr	Leu 1470		Ala
20	Pro	Tyr	Val 147		Val	Tyr	Leu	Leu 178	Asp 0	Asn	Gly	Val	Cys 148		Ala	Lys
	Lys	Lys 149		Lys	Val	Ala	Arg 149		Thr	Leu	Glu	Pro 150		Tyr	Gln	Gln
25	Leu 150		Ser	Phe	Glu	Glu 151		Pro	Gln	Gly	Arg 151		Leu	Gln	Ile	Ile 1520
30	Val	Trp	Gly	Asp	Tyr 152		Arg	Met	Asp	His 153		Ser	Phe	Met	Gly 153	
	Ala	ı Glı	ı Ile	: Leu 154		ı Asp	Glu	Leu	ı Glu 154		Ser	Asn	Met	. Val 155		Gly

Trp Phe Lys Leu Phe Pro Pro Ser Ser Leu Val Asp Pro Thr Ser Ala

Pro Leu Thr Arg Arg Ala Ser Gln Ser Ser Leu Glu Ser Ser Thr Gly Pro Ser Tyr Ser Arg Ser <210> 2 <211> 4980 <212> DNA <213> Mus musculus <400>2gcttccctag ggtggttcgg ctccgccaaa c atg tcg gct ccg ctc ggg ccc Met Ser Ala Pro Leu Gly Pro cgg ggc cgc ccg gct ccc acc ccg gcg gcc tct caa cct cct ccg cag Arg Gly Arg Pro Ala Pro Thr Pro Ala Ala Ser Gln Pro Pro Pro Gln ccc gag atg ccg gac ctc agc cac ctc acg gaa gag gag agg aaa atc Pro Glu Met Pro Asp Leu Ser His Leu Thr Glu Glu Glu Arg Lys Ile atc ctg gct gtc atg gat cgt cag aag aaa gaa gag gag aag gag cag lle Leu Ala Val Met Asp Arg Gln Lys Lys Glu Glu Glu Lys Glu Gln tcc gtg ctc aag atc aaa gaa gaa cac aaa gca caa ccg aca cag tgg Ser Val Leu Lys Ile Lys Glu Glu His Lys Ala Gln Pro Thr Gln Trp

	t t t	cec	t t t	a or t	σσσ	atc	ac t	gaa	ctg	gta	aat	aac	g††	ctg	cag	CCC	2	92
							Thr										_	-
	rne	F10	rne		GIY	116	1111	Ulu		vai	поп	поп	141		UIII	110		
				75					80					85				
5																	0	4.0
							gag										3	40
	Gln	Gln	Lys	Gln	Pro	Asn	Glu	Lys	Glu	Pro	Gln	Thr	Lys	Leu	His	Gln		
			90					95					100					
10	caa	ttt	gaa	atg	tat	aag	gag	caa	gtc	aag	aag	atg	gga	gag	gaa	tcg	3	88
	Gln	Phe	Glu	Met	Tyr	Lys	Glu	Gln	Val	Lys	Lys	Met	Gly	Glu	Glu	Ser		
		105					110					115						
	cag	cag	cag	caa	gag	cag	aag	ggt	gat	gcc	ccg	acc	tgt	ggc	atc	tgc	4	36
15	Gln	Gln	Gln	Gln	Glu	Gln	Lys	Gly	Asp	Ala	Pro	Thr	Cys	Gly	Ile	Cys		
	120					125					130					135		
	cac	aag	aca	aaa	ttt	gca	gat	gga	tgc	ggc	cat	aat	tgt	tcc	tat	tgc	4	84
							Asp											
20		2,0			140		_			145					150			
	caa	acc	ลลซ	ttc	tgt	gc t	cga	tgt	gga	ggt	cga	gtg	tct	tta	cgc	tca	5	32
							Arg											
	UIII	1111	Дуо	155		1114	0	0,0	160		0			165				
95				100					100									
25		0.00	e est t	o t o	taa	ata	tat	aat	tta	toc	രമ	ลลล	caa	caa	ฮลล	atc		580
							Cys											
	ASII	Lys			111	Val	CyS			СУЗ	ni s	Гуз	180		Ulu	110		
			170					175					100					
										1	1		. 4 . 4			a t c		328
30																ctg	,	J 2 0
	Leu			Ser	Gly	Ala			lyr	Asn	Ser			Asn	ınr	Leu		
		185	j				190					195						
	cag	g caa	cct	gat	caa	aag	gtt	cct	cga	ggg	ctt	. cga	aat	gag	gaa	gcc		676

	Gln	Gln	Pro	Asp	Gln	Lys	Val	Pro	Arg	Gly	Leu	Arg	Asn	Glu	Glu	Ala	
	200					205					210					215	
	cct	cag	gag	aag	aaa	gca	aaa	cta	cac	gag	cag	ссс	cag	ttc	caa	gga	724
5											Gln						
9	110	UIII	oru	LJU	220	ma	1 ,5	Deu	1115	225	UIII	110	GIII	1110	230	GI,	
					440					223					200		
	~~~		a a t		+ + 0	+00	at o	a a t	60.0	cr + +	or o or	0.00	g g a	agn	co t	ant	779
											gag						772
	Ala	Pro	Gly		Leu	ser	vai	PTO		vai	Glu	Lys	ч		Ala	1115	
10				235					240					245			
											gga						820
	Gly	Leu	Thr	Arg	Gln	Asp	Thr	He	Lys	Asn	Gly	Ser	Gly	Val	Lys	His	
			250					255					260				
15																	
	cag	att	gcc	agt	gac	atg	cct	tca	gac	aga	aaa	cga	agt	cca	tca	gtg	868
	Gln	Ile	Ala	Ser	Asp	Me t	Pro	Ser	Asp	Arg	Lys	Arg	Ser	Pro	Ser	Val	
		265					270					275					
20	tcc	agg	gat	caa	aat	cga	aga	tac	gag	caa	agt	gaa	gaa	aga	gag	gac	916
	Ser	Arg	Asp	Gln	Asn	Arg	Arg	Tyr	Glu	Gln	Ser	Glu	Glu	Arg	Glu	Asp	
	000																
	280					285					290					295	
	280					285					290					295	
		tca	cag	tat	gtt		tca	gat	ggt	aca	290 atg	cca	aga	tct	cct		964
25	tac		-		_	cct		_	_		atg					tcg	964
25	tac		-		_	cct		_	_							tcg	964
25	tac		-		Val	cct		_	_	Thr	atg				Pro	tcg	964
25	tac Tyr	Ser	Gln	Tyr	Val 300	cct Pro	Ser	Asp	Gly	Thr 305	atg Met	Pro	Arg	Ser	Pro 310	tcg Ser	
25	tac Tyr gat	Ser	Gln	Tyr	Val 300	cct Pro	Ser	Asp	Gly	Thr 305 gag	atg Met	Pro caa	Arg	Ser tat	Pro 310 gaa	tcg Ser	964 1012
	tac Tyr gat	Ser	Gln	Tyr gat Asp	Val 300	cct Pro	Ser	Asp	Gly cgt Arg	Thr 305 gag	atg Met	Pro caa	Arg	Ser tat Tyr	Pro 310 gaa	tcg Ser	
25	tac Tyr gat	Ser	Gln	Tyr	Val 300	cct Pro	Ser	Asp	Gly	Thr 305 gag	atg Met	Pro caa	Arg	Ser tat	Pro 310 gaa	tcg Ser	
	tac Tyr gat Asp	Ser tat Tyr	Gln gct Ala	Tyr gat Asp 315	Val 300 aga Arg	cct Pro cga Arg	Ser tct Ser	Asp cag Gln	Gly cgt Arg 320	Thr 305 gag Glu	atg Met cct Pro	Pro caa GIn	Arg ttt Phe	Ser tat Tyr 325	Pro 310 gaa Glu	tcg Ser gaa Glu	1012
	tac Tyr gat Asp	Ser tat Tyr	Gln gct Ala	Tyr gat Asp 315	Val 300 aga Arg	cct Pro cga Arg	ser tct ser	Asp cag Gln	Gly cgt Arg 320	Thr 305 gag Glu aac	atg Met cct Pro	Pro caa Gln aga	Arg ttt Phe	tat Tyr 325	Pro 310 gaa Glu aga	tcg Ser gaa Glu	
	tac Tyr gat Asp	Ser tat Tyr	Gln gct Ala	Tyr gat Asp 315	Val 300 aga Arg	cct Pro cga Arg	ser tct ser	Asp cag Gln	Gly cgt Arg 320	Thr 305 gag Glu aac	atg Met cct Pro	Pro caa Gln aga	Arg ttt Phe	tat Tyr 325	Pro 310 gaa Glu aga	tcg Ser gaa Glu	1012

	tcc	aaa	gag	tat	att	gtg	gat	gat	gaa	gat	gtg	gag	agc	aga	gat	gaa	1108
				Tyr													
		345					350					355					
5																	
	tat	gaa	aga	caa	agg	aga	gag	gag	gaa	tac	cag	gca	cgc	tac	aga	agt	1156
	Tyr	Glu	Arg	Gln	Arg	Arg	Glu	Glu	Glu	Tyr	Gln	Ala	Arg	Tyr	Arg	Ser	
	360					365					370					375	
10	gat	cca	aat	ctg	gcc	cgg	tat	ccc	gta	aag	cca	caa	ccc	tac	gaa	gaa	1204
	Asp	Pro	Asn	Leu	Ala	Arg	Tyr	Pro	Val	Lys	Pro	Gln	Pro	Tyr		Glu	
					380					385					390		
																	1050
				atc													1252
15	Gln	Met	Arg	He	His	Ala	Glu	Val		Arg	Ala	Arg	HIS		Arg	Arg	
				395					400					405			
					1-1	***	<b></b>		cro t	ara o	a t a	an n	an t	tee	മനന	att	1300
				gtt Val													1000
90	HIS	ser	410		ser	Leu	Ма	415		Olu	Lcu	Gru	420		1110	110	
20			410					710					120				
	tet	ctg	cta	ឧឧଣ	atg	gat	aga	сса	tca	agg	caa	aga	tct	gta	tct	gaa	1348
																Glu	
	501	425				_	430					435					
25																	
	cgt	aga	ı gct	gca	ate	g gaa	aac	caa	. cga	tce	tat	tca	ate	gaa	aga	act	1396
	Arg	g Arg	g Ala	a Ala	Met	Glu	Asn	Gln	Arg	Ser	Туі	Ser	Met	Glu	Arg	Thr	
	440	)				445	;				45(	)				455	
30	cga	a gaş	g gc	t cas	g gga	a caa	agt	tct	tat	cca	a caa	a agg	g acc	tca	ı aat	cat	1444
	Arg	g Glu	u Ala	a Glr	ı Gly	y Gli	ı Sei	Sei	Туг	Pro	Gli	n Arg	g Thi	Sei	Asr	His	
					460	0				46	5				470	)	
	ag	t cc	t cc	c aco	c cc	t cg	g cgs	g age	c cc	t ata	a cc	g ct	t ga	t aga	а сса	a gac	1492

	Ser	Pro	Pro	Thr 475	Pro	Arg	Arg	Ser	Pro 480	Ile	Pro	Leu	Asp	Arg 485	Pro	Asp	
5	_								aaa Lys								1540
10									gaa Glu								1588
15									tcc Ser								1636
10									gga Gly							-	1684
20									gca Ala 560								1732
25									gag Glu								1780
30			Lys						agt Ser						_		1828
		Asn							aag Lys								1876

ı s

	cac	tct	ttg	gaa	gag	gat	ttg	gaa	tgg	tct	gag	cct	cag	att	aag	gac	1924
	His	Ser	Leu	Glu	Glu	Asp	Leu	Glu	Trp	Ser	Glu	Pro	Gln	Ile	Lys	Asp	
					620					625					630		
5																	
	tct	ggg	gta	gat	acc	tgt	agt	agc	aca	acc	ctt	aac	gag	gag	cat	agc	1972
	Ser	Gly	Val	Asp	Thr	Cys	Ser	Ser	Thr	Thr	Leu	Asn	Glu	Glu	His	Ser	
				635					640					645			
10	cat	agt	gat	aag	cac	cct	gtg	acc	tgg	cag	cca	tcc	aaa	gat	gga	gat	2020
	His	Ser	Asp	Lys	His	Pro	Val	Thr	Trp	Gln	Pro	Ser	Lys	Asp	Gly	Asp	
			650					655					660				
											cgt						2068
15	Arg		He	Gly	Arg	He		Leu	Asn	Lys	Arg		Lys	Asp	Gly	Ser	
		665					670					675					
																	9110
	-										tta						2116
90	680	PTO	Arg	ASP	ser	685	Ala	меι	Leu	Gly	Leu 690	Lys	vai	vai	GIY	695	
20	000					000					090					030	
	ลลต	atø	act	gaa	tca	ggt	cga	ctt	tgt	gca	ttt	att	acc	aaa	gta	ааа	2164
											Phe						
	2,0				700	,				705				•	710		
25																	
	aaa	gga	agt	tta	gct	gat	ac t	gta	gga	cat	ctt	aga	cca	ggt	gat	gaa	2212
	Lys	Gly	Ser	Leu	Ala	Asp	Thr	Val	Gly	His	Leu	Arg	Pro	Gly	Asp	Glu	
				715					720					725			
30	gtc	ttg	gaa	tgg	aat	ggg	agg	cta	ttg	caa	gga	gcc	aca	ttt	gag	gaa	2260
	Val	Leu	Glu	Trp	Asn	Gly	Arg	Leu	Leu	Gln	Gly	Ala	Thr	Phe	Glu	Glu	
			730					735					740				
	gtt	tac	aac	att	att	cta	gaa	tcc	aaa	cct	gaa	cca	caa	gtt	gag	ctt	2308

	Val		Asn	Ile	He	Leu		Ser	Lys	Pro	Glu		Gln	Val	Glu	Leu	
		745					750					755					
	gtt	gtt	tca	agg	cca	att	gga	gat	att	cct	aga	ata	cct	gat	agc	acg	2356
5	Val	Val	Ser	Arg	Pro	He	Gly	Asp	He	Pro	Arg	Ile	Pro	Asp	Ser	Thr	
	760					765					770					775	
	ant	go.	000	o t a	aro o	taa	o & t	tot	ngo	tan	+++	an o	tot	000	000	n t cr	2404
									agc								4404
10	nis	Ala	GIII	Leu		261	ser	ser	Ser		rne	GIU	sei	GIII		меι	
10					780					785					790		
	gac	cgt	cct	tct	ata	tcc	gtt	acc	tca	ccc	atg	agt	cct	ggc	atg	ctg	2452
	Asp	Arg	Pro	Ser	Ile	Ser	Val	Thr	Ser	Pro	Met	Ser	Pro	Gly	Met	Leu	
				795					800					805			
15																	
	agg	gat	gtc	ccg	cag	ttc	tta	tct	gga	cag	ctt	tca	ata	aaa	cta	tgg	2500
	Arg	Asp	Val	Pro	Gln	Phe	Leu	Ser	Gly	Gln	Leu	Ser	Ile	Lys	Leu	Trp	
			810					815					820				
20	ttt	gac	aag	gtt	ggt	cac	cag	ttg	ata	gtt	aca	att	ttg	gga	gca	aag	2548
	Phe	Asp	Lys	Val	Gly	His	Gln	Leu	Ile	Val	Thr	Ile	Leu	Gly	Ala	Lys	
		825					830					835					
	gat	ctc	cct	tcc	agg	gaa	gat	ggg	agg	cca	agg	aat	cct	tat	gtt	aag	2596
25	Asp	Leu	Pro	Ser	Arg	Glu	Asp	Gly	Arg	Pro	Arg	Asn	Pro	Tyr	Val	Lys	
	840					845					850					855	
	att	tac	ttc	ctt	cca	gat	aga	agt	gat	aaa	aat	aag	aga	aga	aca	aaa	2644
	He	Tyr	Phe	Leu	Pro	Asp	Arg	Ser	Asp	Lys	Asn	Lys	Arg	Arg	Thr	Lys	
30					860					865					870		
	aca	gtc	aag	aaa	act	ttg	gaa	ccc	aaa	tgg	aac	cag	ac t	ttc	att	tat	2692
	Thr	Val	Lys	Lys	Thr	Leu	Glu	Pro	Lys	Trp	Asn	Gln	Thr	Phe	Ile	Tyr	
				875					880					885			

	tct	cct	gtc	cac	cga	aga	gaa	ttc	cgt	gaa	cga	atg	ctg	gaa	att	acc	2740
	Ser	Pro	Val	His	Arg	Arg	Glu	Phe	Arg	Glu	Arg	Me t	Leu	Glu	He	Thr	
			890					895					900				
5																	
	ctt	tgg	gat	caa	gct	aga	gtt	cga	gaa	gaa	gag	agc	gaa	ttc	tta	gga	2788
	Leu	Trp	Asp	Gln	Ala	Arg	Val	Arg	Glu	Glu	Glu	Ser	Glu	Phe	Leu	Gly	
		905					910					915					
10	gag	att	tta	att	gaa	ttg	gaa	aca	gc t	ttg	cta	gat	gat	gag	ccg	cac	2836
	Glu	He	Leu	Ile	Glu	Leu	Glu	Thr	Ala	Leu	Leu	Asp	Asp	Glu	Pro	His	
	920					925					930					935	
	tgg	tat	aag	ctg	cag	acc	cat	gat	gtc	tcc	tca	ttg	cca	ctc	cct	cgc	2884
15	Trp	Tyr	Lys	Leu	Gln	Thr	His	Asp	Val	Ser	Ser	Leu	Pro	Leu	Pro	Arg	
					940					945					950		
	cct	tcc	cca	tat	ctg	ecc	cgg	agg	cag	ctc	cat	gga	gag	agc	cca	acg	2932
	Pro	Ser	Pro	Tyr	Leu	Pro	Arg	Arg	Gln	Leu	His	Gly	Glu	Ser	Pro	Thr	
20				955					960					965			
	cgc	agg	ctg	caa	agg	tcg	aag	aga	ata	agt	gac	agt	gaa	gtg	tct	gac	2980
	Arg	Arg	Leu	Gln	Arg	Ser	Lys	Arg	He	Ser	Asp	Ser	Glu	Val	Ser	Asp	
			970					975					980				
25																	
	tac	gac	tgc	gag	gat	ggc	gtg	gga	gta	gtg	tca	gat	tat	cga	cac	aat	3028
	Tyr	Asp	Cys	Glu	Asp	Gly	Val	Gly	Val	Val	Ser	Asp	Tyr	Arg	His	Asn	
		985					990					995					
30	ggc	cgc	gat	ctt	caa	agc	tcc	acg	ttg	tcg	gtg	cca	gaa	caa	gtc	atg	3076
	Gly	Arg	Asp	Leu	Gln	Ser	Ser	Thr	Leu	Ser	Val	Pro	Glu	Gln	Val	Met	
	1000	)				1009	5				1010	)				1015	
	tca	tca	aat	cat	tgc	tca	cca	tca	ggg	tct	cct	cat	cga	gta	gat	gtt	3124

	Ser	Ser	Asn	His			Pro	Ser	Gly			His	Arg	Val			
					1020	)				1029	0				1030	)	
	ata	gga	agg	aca	agg	tca	tgg	tcg	cct	agt	gcc	cct	cct	cct	caa	agg	3172
5	Ile	Gly	Arg	Thr	Arg	Ser	Trp	Ser	Pro	Ser	Ala	Pro	Pro	Pro	Gln	Arg	
				1035	õ				1040	)				1045	5		
	aat	gtg	gaa	cag	ggg	cac	cga	ggg	aca	cgt	gc t	ac t	ggc	cat	tac	aac	3220
	Asn	Val	Glu	Gln	Gly	His	Arg	Gly	Thr	Arg	Ala	Thr	Gly	His	Tyr	Asn	
10			1050	)				1055	)				1060	)			
	aca	att	agc	cga	atg	gat	aga	cac	cgt	gtc	atg	gat	gac	cac	tac	tct	3268
	Thr	Ile	Ser	Arg	Met	Asp	Arg	His	Arg	Val	Met	Asp	Asp	His	Tyr	Ser	
		1068	5				1070	)				1075	j				
15																	
	tca	gat	aga	gac	agg	gat	tgt	gaa	gca	gca	gat	aga	cag	cca	tat	cac	3316
	Ser	Asp	Arg	Asp	Arg	Asp	Cys	Glu	Ala	Ala	Asp	Arg	Gln	Pro	Tyr	His	
	1080	)				1085	5				1090	}				1095	
20											cta.						3364
	Arg	Cor	Arc	Sar	Thr	Glu	Gln	Arg	Pro	Leu	1011	Glu	Arg	Ihr	Thr	Thr	
	111 0	201	MIG	261													
	111 6	261	MIS	261	1100	)				1109					1110		
					1100		cgt	cct	gat	1109	5	ctc	atg	agg		)	3412
25	cgc	tcc	aga	tcc	1100 tct	gaa				1109 aca	aac				tcg	) atg	3412
25	cgc	tcc	aga	tcc Ser	tct Ser	gaa				1109 aca Thr	5				tcg Ser	) atg	3412
25	cgc	tcc	aga	tcc	tct Ser	gaa			Asp	1109 aca Thr	aac			Arg	tcg Ser	) atg	3412
25	cgc Arg	tcc Ser	aga Arg	tcc Ser	tct Ser	gaa Glu	Arg	Pro	Asp 1120	1109 aca Thr	aac	Leu	Met	Arg 1125	tcg Ser	atg Met	3412 3460
25	cgc Arg	tcc Ser tca	aga Arg tta	tcc Ser 1115	tct Ser 5	gaa Glu gga	Arg aga	Pro tct	Asp 1120 gcc	aca Thr	aac Asn	Leu tca	Met	Arg 1125 gcc	tcg Ser	atg Met tcg	
25	cgc Arg	tcc Ser tca	aga Arg tta	tcc Ser 1115 atg	tct Ser 5	gaa Glu gga	Arg aga	Pro tct	Asp 1120 gcc Ala	aca Thr	aac Asn	Leu tca	Met	Arg 1125 gcc Ala	tcg Ser	atg Met tcg	
	cgc Arg	tcc Ser tca	aga Arg tta Leu	tcc Ser 1115 atg	tct Ser 5	gaa Glu gga	Arg aga	Pro tct Ser	Asp 1120 gcc Ala	aca Thr	aac Asn	Leu tca	Met cct Pro	Arg 1125 gcc Ala	tcg Ser	atg Met tcg	
	cgc Arg cct Pro	tcc Ser tca Ser	aga Arg tta Leu 113	tcc Ser 1115 atg Met	tct Ser 5 act	gaa Glu gga Gly	Arg aga Arg	tct Ser 1135	Asp 1120 gcc Ala	aca Thr ) cct Pro	aac Asn	Leu tca Ser	cct Pro	Arg 1125 gcc Ala	tcg Ser tta Leu	atg Met tcg Ser	
	cgc Arg cct Pro	tcc Ser tca Ser	aga Arg tta Leu 1130	tcc Ser 1115 atg Met O	tct Ser 5 act Thr	gaa Glu gga Gly	Arg aga Arg	tct Ser 1135	Asp 1120 gcc Ala	aca Thr  cct Pro	aac Asn cct Pro	tca Ser	cct Pro 1140	Ars 1125 gcc Ala )	tcg Ser 5 tta Leu	atg Met tcg Ser	3460

	ccg	gga	aca	gga	cga	agg	ggc	cga	cag	ctt	cca	cag	ctt	cca	cca	aag	3556
	Pro	Gly	Thr	Gly	Arg	Arg	Gly	Arg	Gln	Leu	Pro	Gln	Leu	Pro	Pro	Lys	
	1160	)				1165	)				1170	)				1175	
5																	
	gga	aca	ttg	gag	aga	agt	gc t	atg	gat	ata	gag	gag	aga	aat	cgc	caa	3604
	Gly	Thr	Leu	Glu	Arg	Ser	Ala	Met	Asp	He	Glu	Glu	Arg	Asn	Arg	Gln	
					1180	)				1185	<u>,</u>				1190	)	
10	atg	aaa	ctt	aac	aaa	tac	aaa	cag	gta	gcc	gga	tca	gac	ccc	aga	ctg	3652
	Met	Lys	Leu	Asn	Lys	Tyr	Lys	Gln	Val	Ala	Gly	Ser	Asp	Pro	Arg	Leu	
				119	5				1200	}				1205	5		
	gag	caa	gat	tac	cat	tcg	aag	tat	cgc	tca	gga	tgg	gat	cca	cat	aga	3700
15	Glu	Gln	Asp	Tyr	His	Ser	Lys	Tyr	Arg	Ser	Gly	Trp	Asp	Pro	His	Arg	
			1210	)				1215	j .				1220	)			
	ggg	gca	gat	ac t	gtt	tcc	ac t	aaa	tcc	tcg	gac	agt	gat	gta	agt	gat	3748
	Gly	Ala	Asp	Thr	Val	Ser	Thr	Lys	Ser	Ser	Asp	Ser	Asp	Val	Ser	Asp	
20		1225	5				1230	)				1235	5				
	gta	tct	gcg	gtt	tca	agg	ac t	agt	agt	gct	tct	cgt	ttc	agc	agc	aca	3796
				_													
	Val	Ser		-	Ser	Arg	Thr	Ser	Ser	Ala		Arg	Phe	Ser	Ser	Thr	
	Val 1240			-	Ser	Arg 1249		Ser	Ser	Ala			Phe	Ser	Ser	Thr 1255	
25				-	Ser			Ser	Ser	Ala	Ser		Phe	Ser	Ser		
25	1240	)	Ala	Val		124	ō		Ser		Ser 1250	)				1255	3844
25	1240 agc	) tac	Ala	Val	gtc	124s	tca	gag		ccg	Ser 1250 aga	) gga	aac	agg	aaa	1255 atc	
25	1240 agc	) tac	Ala	Val	gtc	1245 caa Gln	tca	gag	cgg	ccg	Ser 1250 aga Arg	) gga	aac	agg	aaa	1255 atc Ile	
25	1240 agc	) tac	Ala	Val	gtc Val	1245 caa Gln	tca	gag	cgg	ccg Pro	Ser 1250 aga Arg	) gga	aac	agg	aaa Lys	1255 atc Ile	
25	agc Ser	tac Tyr	Ala atg Met	Val tcc Ser	gtc Val 126	taa caa Gln 0	tca Ser	gag Glu	cgg	ccg Pro 126	Ser 1250 aga Arg	gga Gly	aac Asn	agg Arg	aaa Lys 1270	1255 atc Ile	
	agc Ser	tac Tyr	Ala atg Met	Val tcc Ser	gtc Val 1260	caa Gln 0	tca Ser	gag Glu caa	cgg Arg	ccg Pro 126	Ser 1250 aga Arg	gga Gly atg	aac Asn	agg Arg	aaa Lys 1270	atc Ile )	3844
	agc Ser	tac Tyr	Ala atg Met	Val tcc Ser	gtc Val 126 tcc Ser	caa Gln 0	tca Ser	gag Glu caa	cgg Arg	ccg Pro 1269 aga Arg	Ser 1250 aga Arg	gga Gly atg	aac Asn	agg Arg	aaa Lys 1270 tcg Ser	atc Ile )	3844
	agc Ser	tac Tyr	Ala atg Met	tcc Ser aca	gtc Val 126 tcc Ser	caa Gln 0	tca Ser	gag Glu caa	cgg Arg aac Asn	ccg Pro 1269 aga Arg	Ser 1250 aga Arg	gga Gly atg	aac Asn	agg Arg gtg Val	aaa Lys 1270 tcg Ser	atc Ile )	3844

	Lys	Asn	Leu	Thr	Lys	Ser	Thr	Ser	He	Ser	Gly	Asp	Met	Cys	Ser	Leu	
			1290	)				1295	)				1300	)			
											gca						3988
5	Glu	Lys	Asn	Asp	Gly	Ser	Gln	Ser	Asp	Thr	Ala	Val	Gly	Ala	Leu	Gly	
		1305	ĵ				1310	)				1315	)				
	acc	agt	ggc	aag	aag	cgg	cga	tct	agc	att	ggg	gcc	aaa	atg	gta	gc t	4036
	Thr	Ser	Gly	Lys	Lys	Arg	Arg	Ser	Ser	He	Gly	Ala	Lys	Me t	Val	Ala	
10	1320	)				1325	5				1330	)				1335	
	att	gtt	ggt	ctc	tca	cgg	aaa	agt	cgc	agt	gcc	tct	caa	ctc	agc	caa	4084
	Ile	Val	Gly	Leu	Ser	Arg	Lys	Ser	Arg	Ser	Ala	Ser	Gln	Leu	Ser	Gln	
					1340	)				1345	)				1350	)	
15																	
	acc	gaa	gga	gga	ggt	aaa	aag	cta	cgg	agc	ac t	gtt	cag	aga	agc	acg	4132
	Thr	Glu	Gly	Gly	Gly	Lys	Lys	Leu	Arg	Ser	Thr	Val	Gln	Arg	Ser	Thr	
				1359	5				1360	)				1365	5		
20	gag	acc	ggg	cta	gca	gtg	gag	atg	agg	aac	tgg	atg	acc	cgc	cag	gcc	4180
	Glu	Thr	Gly	Leu	Ala	Val	Glu	Met	Arg	Asn	Trp	Met	Thr	Arg	Gln	Ala	
			137	0				1375	5				1386	)			
	agc	cgg	gaa	tcc	aca	gat	ggc	agc	atg	aac	agc	tat	agc	tcg	gaa	gga	4228
25	Ser	Arg	Glu	Ser	Thr	Asp	Gly	Ser	Met	Asn	Ser	Tyr	Ser	Ser	Glu	Gly	
		138	5				1390	)				1395	5				
	aat	ctg	atc	ttc	cct	ggg	gtc	cgc	ctg	gcc	tct	gac	agc	cag	ttc	agt	4276
	Asn	Leu	He	Phe	Pro	Gly	Val	Arg	Leu	Ala	Ser	Asp	Ser	Gln	Phe	Ser	
30	140	0				140	5				1410	)				1415	
																	4904
	gat	ttc	ctg	gat	ggc	ctg	ggc	cct	gct	cag	cta	gtg	gga	cgc	cag	acc	4324
											cta Leu						4324

	ctg	gct	ac t	cct	gca	atg	ggt	gac	att	cag	gtg	gga	atg	atg	gat	aaa	4372
	Leu	Ala	Thr	Pro	Ala	Met	Gly	Asp	He	Gln	Val	Gly	Me t	Me t	Asp	Lys	
				143	5				1440	)				144	5		
5																	
	aag	gga	cag	ctg	gag	gta	gaa	atc	atc	cgg	gcg	cgc	ggc	ctt	gtg	gta	4420
	Lys	Gly	Gln	Leu	Glu	Val	Glu	He	He	Arg	Ala	Arg	Gly	Leu	Val	Val	
			1450	)				1455	5				1460	)			
10	aaa	cca	ggt	tcc	aag	aca	ctg	cca	gca	ccg	tat	gtc	aag	gtg	tat	ctg	4468
	Lys	Pro	Gly	Ser	Lys	Thr	Leu	Pro	Ala	Pro	Tyr	Val	Lys	Val	Tyr	Leu	
		1465	5				1470	)				1475	5				
	tta	gac	aac	gga	gtc	tgc	ata	gcc	aaa	aag	aaa	acc	aag	gtg	gcg	aga	4516
15	Leu	Asp	Asn	Gly	Val	Cys	He	Ala	Lys	Lys	Lys	Thr	Lys	Val	Ala	Arg	
	1780	)				1485	)				1490	)				1495	
	aag	acc	ctg	gag	ccc	ctg	tac	cag	cag	ctc	ttg	tcc	ttc	gag	gag	agc	4564
	Lys	Thr	Leu	Glu	Pro	Leu	Tyr	Gln	Gln	Leu	Leu	Ser	Phe	Glu	Glu	Ser	
20					1500	)				1505	<u>,                                    </u>				1510	)	
	ccc	cag	ggg	agg	gtg	tta	cag	atc	att	gtc	tgg	gga	gat	tat	ggt	cgt	4612
	Pro	Gln	Gly	Arg	Val	Leu	Gln	He	He	Val	Trp	Gly	Asp	Tyr	Gly	Arg	
				1515	5				1520	}				1525	)		
25																	
	atg	gat	cac	aaa	tcc	ttt	atg	gga	gtg	gcc	cag	ata	ctc	tta	gat	gaa	4660
	Met	Asp	His	Lys	Ser	Phe	Met	Gly	Val	Ala	Gln	Ile	Leu	Leu	Asp	Glu	
			1530	)				1535	5				1540	)			
30	ctg	gaa	cta	tcc	aac	atg	gtg	att	gga	tgg	ttc	aaa	ctc	ttc	cct	cct	4708
	Leu	Glu	Leu	Ser	Asn	Me t	Val	He	Gly	Trp	Phe	Lys	Leu	Phe	Pro	Pro	
		1548	5				1550	)				1555	5				
	tcc	tcc	cta	gta	gat	cca	acc	tcg	gca	cct	ctg	aca	aga	aga	gct	tcc	4756

		Pro Thr Ser Ala Pro 1565	Leu Thr Arg Arg Ala Ser 1570 1575	
5	caa tcg tct ctg gaa a Gln Ser Ser Leu Glu S 1580		Ser Tyr Ser Arg Ser	4801
	tagcaactat aaaactgttg	g tcacaacaac cagcga	taca aaaaccagaa gaaaacgcac	4861
10	aggtggaagc ccctggtaac	c actgcatgct tgatgt	tgtg tctacagagc ccacgtctag	4921
	ggataccaag cagtcctgts	g ttctcagagg aagtcg	taca cattgtgccc tagcaaagg	4980
15	<210> 3 <211> 45 <212> DNA <213> Mus musculus			
20	<400> 3 ttgcgctcac tcttctggcc	c tecettgeca ttetge	tctg aaagc	45
	<210> 4 <211> 16 <212> DNA			
25	<213> Mus musculus			
	<400> 4 acctacgtga ctacgt			16
30	<210> 5 <211> 16 <212> PRT <213> Mus musculus			

<400> 5

Gln Met Ser His Arg Leu Glu Pro Arg Arg Pro

1

5

10

#### WHAT IS CLAIMED IS:

- 1. A protein having the amino acid sequence set forth under SEQ ID NO:1 in the Sequence Listing.
- 2. A protein having an amino acid sequence with one or more amino acids deleted, substituted, inserted or added relative to the amino acid sequence set forth under SEQ ID NO:1 in the Sequence Listing and which has a property to interact with GDP/GTP exchange factor II.
  - 3. A mouse gene which encodes the protein of claim 1 or 2.
- 4. A DNA having a nucleotide sequence set forth under SEQ ID NO:2 in the Sequence Listing, the DNA being a cDNA corresponding to the protein of claim 1.
- 5. A DNA having a nucleotide sequence with one or more nucleotides deleted, substituted, inserted or added relative to the nucleotide sequence set forth under SEQ ID NO:2 in the Sequence Listing and encoding the protein of claim 1 or 2.
- 6. A DNA having the nucleotide sequence of the coding region of the DNA of claim 4.
- 7. A DNA having the nucleotide sequence of the coding region of the DNA of claim 5.
  - 8. A DNA fragment consisting of a part of the DNA of claim 4.
  - 9. A probe comprising a DNA which hybridizes with the DNA of claim 4.
- 10. A primer DNA fragment consisting of a partial sequence of the sequence of one of claims 4 to 7.
  - 11. A recombinant vector having the DNA of claim 4.
  - 12. A recombinant vector having the DNA of claim 5.
- 13. A monoclonal or polyclonal antibody directed to the protein of claim 1 or 2.
- 14. A diagnostic agent for secretory disorders or brain-nervous system diseases comprising the probe of claim 9.
- 15. A diagnostic agent for secretory disorders or brain-nervous system diseases comprising the antibody of claim 13.

5

### ABSTRACT

Provided is a protein used in the development of a therapeutic agent for neuron- or endocrine cell-related diseases, in which the transport system is involved. The protein has an amino acid sequence with one or more amino acids deleted, substituted, inserted or added relative to the amino acid sequence set forth under SEQ ID NO:1 in the Sequence Listing and which has a property to interact with GDP/GTP exchange factor II.

# Figure 1

130 439 218 342 DIGTNWYAVLA SLDVKVSETSSHQDAVTICTLGIGTAF SIL-DNTP EEGTSWYIILK SVNV-VIYGKG----V-VCTLHEGDDF KLALVNDAP DEGDNFYVIDQ EMDVYVNNEWAT----SVGEGGSF LALIYGTP EPGDEFFIILE TAAV-LQRRSENEEFVEVGRLGPSDYF HIALLMNRP CAMP-A CAMP-B RI a-A RI a-B

Figure 2




Figure 3




Figure 4

	CO	S-1		MI	N6		Br	ain
-	GST-Rim2	GST	-	GST-Rim2	GST	_	GST-Rim1	GS-
FLAG- cAMP-GEFII			cAMP-GEFII	-		cAMP-GEFII		

Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12





Figure 13



# Declaration and Power of Attorney For Utility or Design Patent Application 特許出願宣言書

## Japanese Language Declaration

私は、下欄に氏名を記載した発明 宣言する:	明者として、以下のとお	; <b>n</b>	As a below named inventor, I hereby declare	e that:				
私の住所、郵便の宛先および国籍は であり、	は、下欄に氏名に続いて記	己載したとおり	My residence, post office address and citiz below next to my name.	zenship are as stated				
名称の発明に関し、請求の範囲に 最初にして唯一の発明者である(一 場合)か、もしくは本来の、最初に「 下欄に記載されている場合)と信じ	人の氏名のみが下欄に記 して共同の発明者である	己載されている	I believe I am the original, first and sole inventisted below) or an original, first and joint invare listed below) of the subject matter which a patent is sought on the invention en	ventor (if plural names ch is claimed and for				
			Protein Rim2					
その明細書を			the specification of which					
。。。。。。。。。。。。。。。。。。。。。。。。。。。。。。。。。。。。			(check one) ☑ is attached hereto.					
<u> </u>		日に出願番号	was filed on					
皇 第			Application No.					
	F	に補正した。	and was amended on					
(該当する場合)	)			f applicable)				
上で、私は、前記のとおり補正した請求 は、理解したことを陳述する。 は、連邦規則法典第37部第1章 情報を開示すべき義務を有すること がは合衆国法典第35部第119条(a・ 記の外国特許出願又は発明者証出願	章第56条に従い、本題の とを認める。 一d)項又は第365条(b)項	)審査に所要の 質に基づく、下	I hereby state that I have reviewed and under the above identified specification, including the by any amendment referred to above. I acknowledge the duty to disclose information the examination of this application in accordance of Federal Regulations, §1.56.	e claims, as amended on which is material to nce with Title 37, Code				
くても米国以外の1ヶ国を指名した」 し、更に優先権の主張に係わる基礎 特許出願、又は発明者証出願或るい	PCT国際出願の外国優先 出願の出願日前の出願日	-権利益を主張 ∃を有する外国	I hereby claim foreign priority benefits under Title 35, United Code §119(a-d) or §365(b) of any foreign application(s) for or inventor's certificate, or §365(a) of any PCT internapplication which designated at least one country other the United States of America, listed below and have also idealed below, by checking the "No" box, any foreign application for or inventor's certificate, or of any PCT international application a filing date before that of the application on which priority is clean.					
Prior foreign applications 先の外国出願 288372 / 99	JAPAN	08 / 1	.0 / 1999	Priority claimed 優先権の主張				
	intry)	(Day/Month/Yea (出願の年月日)		VI □ Yes No あり なし				
(Number) (Cou (番号) (国名	intry) i)	(Day/Month/Yea (出願の年月日)	r Filed)	Yes No ab &L				
□ その他の外国特許出願番号は別	別紙の追補優先権欄にて	記載する。	☐ Additional foreign application numb	ers are listed on a				
	The second second		supplemental priority sheet attached hereto.					

## Japanese Language Utility or Design Patent Application Declaration

私は、合衆国法典第35部第119条(e)項に基づく、下記の合衆国仮特許出 願の利益を主張する。

I hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional application(s) listed below.

(Number) (番号)	(Day/Month/Ye 出願の年月日	ear Filed)					
(Number) (番号)	(Day/Month/Ye 出願の年月日	ear Filed)					
(Number) (番号)	(Day/Month/Ye 出願の年月日	ear Filed)					
□ その他の合衆国仮特許出願番号は別総	<b>氏の追補優先権欄にて記載する。</b>	☐ Additional provisional apsupplemental priority sheet a	oplication numbers are listed on a ttached hereto.				
私は、合衆国法典第35部第120条に基づ 第365条(c)項に基づく合衆国を指名したP 願の請求の範囲各項に記載の主題が合衆国 態様で、先の合衆国特許出願又はPCT国際 おいて、先の出願の出願日と本願の国内出 有効となった連邦規則法典第37部第1章 の情報を開示すべき義務を有することを記 (Application No.)	CT国際出願の利益を主張し、本 国法典第35部第112条第1項規定の 民出願に開示されていない限度に 出願日又はPCT国際出願日の間に 第56条に記載の特許要件に所要	I hereby claim the benefit under Title 35, United States Code §12t of any United States application(s), or §365(c) of any PC international application designating the United States of Americal listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PC international application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability at defined in Title 37, Code of Federal Regulations §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.					
(Application No.) (出顧番号)	(Day/Month/Year Filed) (出願の年月日)	(現況) (特許済み、係属中 放棄済み)	(Status) (patented, pending, abandoned)				
== ³ (Application No.) =± ⁸ (出願番号) =±	(Day/Month/Year Filed) (出願の年月日)	(現況) (特許済み、係属中 放棄済み)	(Status) (patented, pending, abandoned)				
 	は別紙の追補優先権欄にて記載	版					
私は、ここに自己の知識にもとずいて行っ己の有する情報および信ずるところに従っじ、さらに故意に虚偽の陳述等を行った場により、罰金もしくは禁錮に処せられるれ、またかかる故意による虚偽による陳述	て行った陳述が真実であると信 8合、合衆国法典第18部第1001条 か、またはこれらの刑が併科さ	knowledge are true and that and belief are believed to be tr were made with the knowledg	atements made herein of my own all statements made on information ue; and further that these statements e that willful false statements and the				

私、下記署名者は、ここに記載の米国弁護士または代理人に本出願に関 し特許商標庁にて取られるいかなる行為に関して、同米国弁護士又は代理 人が、私に直接連絡なしに私の外国弁護士或るいは法人代表者からの指示 を受け取り、それに従うようここに委任する。この指示を出す者が変更の 場合には、ここに記載の米国弁護士又は代理人にその旨通知される。

される特許の有効性を損なうことがあることを認識して、以上の陳述を

行ったことを宣言する。

like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application

or any patent issued thereon.

The undersigned hereby authorizes the U.S. attorney or agent named herein to accept and follow instructions from either his foreign patent agent or corporate representative, if any, as to any action to be taken in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorney or agent and the undersigned. In the event of a change in the persons from whom instructions may be taken, the U.S. attorney or agent named herein will be so notified by the undersigned.

# Japanese Language Utility or Design Patent Application Declaration

委任状: 私は、下記発明者として、下記に明記された顧客番号を伴う以下の弁護士又は、代理人をここに選任し、本順の手続きを遂行すること並びにこれに関する一切の行為を特許商標庁に対して行うことを委任する。そして全ての通信はこの顧客番号宛に発送される。

#### 顧客番号 7055

現在選任された弁護士は下記の通りである。

POWER OF ATTORNEY: As a named inventor, I hereby appoint the attorney(s) and/or agent(s) associated with the Customer Number provided below to prosecute this application and transact all business in the Patent and Trademark Office connected therewith, and direct that all correspondence be addressed to that Customer Number:

#### **CUSTOMER NUMBER 7055**

The appointed attorneys presently include:

Neil F. GreenblumReg. No. 28,394Bruce H. BernsteinReg. No. 29,027Roger P. GlassReg. No. 30,841James L. RowlandReg. No. 32,674Arnold TurkReg. No. 33,094

Address: GREENBLUM & BERNSTEIN, P.L.C.

1941 ROLAND CLARKE PLACE RESTON, VA 20191

直接電話連絡先:(名称および電話番号)

THE WAY

Direct Telephone Calls to: (name and telephone number)

#### GREENBLUM & BERNSTEIN, P.L.C.

(703) 716-1191

・ ・ ・ ・ ・ ・ ・ ・ ・ ・ ・ ・ ・ ・ ・ ・ ・ ・ ・	
唯一のまたは第一の発明者の氏名	Full name of sole or first inventor  Susumu SEINO  June 29, 2000
一、同発明者の署名 日付	Susumu SEINO June 29, 2000 Inventor's signature Date  Susumu Kurs June 29, 200
住所	Residence Japan
五 国籍	Citizenship Japan
郵便の宛先	Post Office Address 22-1-4 Aobanomorinomachi, 638-1 Chibateramachi,
	Chuo-ku, Chiba-shi, Chiba 260-0844 Japan
第2の共同発明者の氏名(該当する場合)	Full name of second joint inventor, if any Tadao SHIBASAKI
同第2共同発明者の署名 川付	Second Inventor's signature Date  Tadao Philasahi Tune 29, 200
住所	Residence Japan
国籍	Citizenship Japan
郵便の宛先	Post Office Address Yamawa mansion 103, 991-128 Yahagi-cho,
	Chuo-ku, Chiba-shi, Chiba 260-0851 Japan

(第六またはそれ以降の共同発明者に対しても同様な情報 および署名を提供すること。)

(Supply similar information and signature for third and subsequent joint inventors.)

Page 3 of 5

# Japanese Language Utility or DesignPatent Application Declaration

第三の共同発明者の氏名	. 10	Full name of third inventor, if any
共同発明者の署名	日付	Nobuaki OZAKI  Third Inventor's signature Jobnaki Gaki June 8,  Residence
住所		
国籍		Japan
		Cıtizenship Japan
郵便の宛先   		Post Office Address Excel-Tsurumai 65, 4-28-1 Yamawaki-cho,
		Showa-ku, Nagoya-shi, Aichi 466-0063 Japan
第四の共同発明者の氏名		Full name of fourth inventor, if any
共同発明者の署名	日付	Fourth Inventor's signature Date
住所		Residence
===国籍 =================================		Citizenship
郵便の宛先		Post Office Address
第五の共同発明者の氏名		Full name of fifth inventor, if any
共同発明者の署名	日付	Fifth Inventor's signature Date
注所		Residence
三		Citizenship
郵便の宛先		Post Office Address
第六の共同発明者の氏名		Full name of sixth inventor, if any
共同発明者の署名	日付	Sixth Inventor's signature Date
住所		Residence
国籍		Citizenship
郵便の宛先		Post Office Address
		the control of the co

(それ以降の共同発明者にたいしても同様な情報 および署名を提供すること。)

(Supply similar information and signature for subsequent joint inventors.)

# United States Patent & Trademark Office Office of Initial Patent Examination -- Scanning Division



for	scanning.		(Document title)	<del></del>	•
	Page(s)	of		were	not presen
	Page(s 500 Seanning)	vere found au of DlC	land From (Document title)	were	not presen
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1					

☐ Scanned copy is best available